

Counting faces of graphical zonotopes

Vladimir Grujić *

University of Belgrade, Faculty of Mathematics,
Studentski trg 16, Belgrade, Serbia

Received 15 June 2016, accepted 16 January 2017, published online 6 March 2017

Abstract

It is a classical fact that the number of vertices of the graphical zonotope Z_Γ is equal to the number of acyclic orientations of a graph Γ . We show that the f -polynomial of Z_Γ is obtained as the principal specialization of the q -analog of the chromatic symmetric function of Γ .

Keywords: Graphical zonotope, f -vector, graphical matroid, symmetric function.

Math. Subj. Class.: 05E05, 52B05, 16T05

1 Introduction

The f -polynomial of an n -dimensional polytope P is defined by $f(P, q) = \sum_{i=0}^n f_i(P)q^i$, where $f_i(P)$ is the number of i -dimensional faces of P . The f -polynomial $f(Z_\Gamma, q)$ of the graphical zonotope Z_Γ is a combinatorial invariant of a finite, simple graph Γ . The vertices of Z_Γ are in one-to-one correspondence with regions of the graphical hyperplane arrangement \mathcal{H}_Γ , which are enumerated by acyclic orientations of Γ .

Stanley's chromatic symmetric function $\Psi(\Gamma) = \sum_{f \text{ proper}} \mathbf{x}_f$ of a graph $\Gamma = (V, E)$, introduced in [7], is the enumerator function of proper colorings $f: V \rightarrow \mathbb{N}$, where $\mathbf{x}_f = x_{f(1)} \cdots x_{f(n)}$ and f is proper if there are no monochromatic edges. The chromatic polynomial $\chi(\Gamma, d)$ of the graph Γ , which counts proper colorings with a finite number of colors, appears as the principal specialization

$$\chi(\Gamma, d) = \mathbf{ps}(\Psi(\Gamma))(d) = \Psi(\Gamma) \big|_{x_1=\cdots=x_d=1, x_{d+1}=\cdots=0}.$$

The number of acyclic orientations of Γ is determined by the value of the chromatic polynomial $\chi(\Gamma, d)$ at $d = -1$, [6]

$$a(\Gamma) = (-1)^{|V|} \chi(\Gamma, -1). \quad (1.1)$$

* Author is supported by Ministry of Education, Science and Technological developments of Republic of Serbia, Project 174034.

E-mail address: vgrujic@matf.bg.ac.rs (Vladimir Grujić)

There is a q -analog of the chromatic symmetric function $\Psi_q(\Gamma)$ introduced in a wider context of the combinatorial Hopf algebra of simplicial complexes considered in [2]. It is a symmetric function over the field of rational functions in q . The principal specialization of $\Psi_q(\Gamma)$ is the q -analog of the chromatic polynomial $\chi_q(\Gamma, d)$.

The main result of this paper is the following generalization of formula (1.1):

Theorem 1.1. *Let $\Gamma = (V, E)$ be a simple connected graph and \mathcal{Z}_Γ the corresponding graphical zonotope. Then the f -polynomial of \mathcal{Z}_Γ is given by*

$$f(\mathcal{Z}_\Gamma, q) = (-1)^{|V|} \chi_{-q}(\Gamma, -1).$$

The cancellation-free formula for the antipode in the Hopf algebra of graphs, obtained by Humpert and Martin in [3], reflects the fact that $f(\mathcal{Z}_\Gamma, q)$ depends only on the graphical matroid $M(\Gamma)$ associated to Γ . For instance, for any tree T_n the graphical matroid is the uniform matroid $M(T_n) = U_n^n$ and the corresponding graphical zonotope is the cube $\mathcal{Z}_{T_n} = I^{n-1}$. Whitney’s theorem from 1933 describes how two graphs with the same graphical matroid are related [9]. It can be used to find more interesting nonisomorphic graphs with the same f -polynomials of corresponding graphical zonotopes.

The paper is organized as follows. In Section 2, we review the basic facts about zonotopes. In Section 3, the q -analog of the chromatic symmetric function $\Psi_q(\Gamma)$ of a graph Γ is introduced. Theorem 1.1 is proved in Section 4. We present some examples and calculations in Section 5.

2 Zonotopes

A zonotope $\mathcal{Z} = \mathcal{Z}(v_1, \dots, v_m)$ is a convex polytope determined by a collection of vectors $\{v_1, \dots, v_m\}$ in \mathbb{R}^n as the Minkowski sum of line segments

$$\mathcal{Z} = [-v_1, v_1] + \dots + [-v_m, v_m].$$

It is a projection of the m -cube $[-1, 1]^m$ under the linear map $\mathbf{t} \mapsto A\mathbf{t}$, $\mathbf{t} \in [-1, 1]^m$, where $A = [v_1 \dots v_m]$ is an $n \times m$ -matrix whose columns are vectors v_1, \dots, v_m . The zonotope \mathcal{Z} is symmetric about the origin and all its faces are translations of zonotopes.

To a collection of vectors $\{v_1, \dots, v_m\}$ is associated a central arrangement of hyperplanes $\mathcal{H} = \{H_{v_1}, \dots, H_{v_m}\}$, where H_v denotes the hyperplane perpendicular to a vector $v \in \mathbb{R}^n$. The zonotope \mathcal{Z} and the corresponding arrangement of hyperplanes \mathcal{H} are closely related. In fact the associated fan $\mathcal{F}_\mathcal{H}$ of the arrangement \mathcal{H} is the normal fan $\mathcal{N}(\mathcal{Z})$ of the zonotope \mathcal{Z} (see [10, Theorem 7.16]). It follows that the face lattice of $\mathcal{F}_\mathcal{H}$ and the reverse face lattice of \mathcal{Z} are isomorphic. In particular, vertices of \mathcal{Z} correspond to regions of \mathcal{H} and their total numbers coincide

$$f_0(\mathcal{Z}) = r(\mathcal{H}). \tag{2.1}$$

The faces of the zonotope \mathcal{Z} are encoded by covectors of the oriented matroid \mathcal{M} associated to the collection of vectors $\{v_1, \dots, v_m\}$. The covectors are sign vectors

$$\mathcal{V}^* = \{\text{sign}(v) \in \{+, -, 0\}^m \mid v \in \mathbb{R}^n\},$$

where $\text{sign}(v)_i = \begin{cases} +, & \langle v, v_i \rangle > 0 \\ 0, & \langle v, v_i \rangle = 0 \\ -, & \langle v, v_i \rangle < 0 \end{cases}$, $i = 1, \dots, m$. The face lattice of the zonotope \mathcal{Z}

is isomorphic to the lattice of covectors componentwise induced by $+, - < 0$ on \mathcal{V}^* .

A special class of zonotopes is determined by simple graphs. To a connected graph $\Gamma = (V, E)$, whose vertices are enumerated by integers $V = \{1, \dots, n\}$, are associated the *graphical zonotope*

$$\mathcal{Z}_\Gamma = \mathcal{Z}(e_i - e_j \mid i < j, \{i, j\} \in E)$$

and the *graphical arrangement* in \mathbb{R}^n

$$\mathcal{H}_\Gamma = \{H_{e_i - e_j} \mid i < j, \{i, j\} \in E\}.$$

There is a bijective correspondence between regions of \mathcal{H}_Γ and acyclic orientations of Γ , [8, Proposition 2.5], which by (2.1) implies

$$f_0(\mathcal{Z}_\Gamma) = r(\mathcal{H}_\Gamma) = a(\Gamma). \tag{2.2}$$

The arrangement \mathcal{H}_Γ is refined by the braid arrangement \mathcal{A}_{n-1} consisting of all hyperplanes $H_{e_i - e_j}, 1 \leq i < j \leq n$. Thus \mathcal{Z}_Γ belongs to a wider class of convex polytopes called generalized permutohedra introduced in [4]. Since arrangements \mathcal{H}_Γ and \mathcal{A}_{n-1} are not essential we take their quotients by the line $l : x_1 = \dots = x_n$ and without confusing retain the same notation. Consequently $\dim \mathcal{Z}_\Gamma = n - 1$.




Figure 1: Permutohedron Pe^3 and cube I^3 .

Example 2.1. (i) The permutohedron Pe^{n-1} is represented as the graphical zonotope \mathcal{Z}_{K_n} corresponding to the complete graph K_n on n vertices (Figure 1).

(ii) The cube I^{n-1} is represented as the graphical zonotope \mathcal{Z}_{T_n} corresponding to an arbitrary tree T_n on n vertices. This shows that the graph Γ is not determined by the combinatorial type of the zonotope \mathcal{Z}_Γ .

3 q -analog of chromatic symmetric function of graph

Stanley’s chromatic symmetric function $\Psi(\Gamma)$ can be obtained in a purely algebraic way. A combinatorial Hopf algebra \mathcal{H} is a graded, connected Hopf algebra equipped with the multiplicative linear functional $\zeta : \mathcal{H} \rightarrow \mathbf{k}$ to the ground field \mathbf{k} . For the theory of combinatorial Hopf algebras see [1]. Consider the combinatorial Hopf algebra of graphs \mathcal{G} which is linearly generated over a field \mathbf{k} by simple finite graphs with the product defined by disjoint union $\Gamma_1 \cdot \Gamma_2 = \Gamma_1 \sqcup \Gamma_2$ and the coproduct

$$\Delta(\Gamma) = \sum_{I \subset V} \Gamma|_I \otimes \Gamma|_{V \setminus I},$$

where $\Gamma|_I$ denotes the induced subgraph on $I \subset V$. The structure of \mathcal{G} is completed by the character $\zeta: \mathcal{G} \rightarrow \mathbf{k}$ defined to be $\zeta(\Gamma) = 1$ for Γ with no edges and $\zeta(\Gamma) = 0$ otherwise. Then it turns out that $\Psi(\Gamma)$ is the image of the unique morphism of combinatorial Hopf algebras to symmetric functions $\Psi: \mathcal{G} \rightarrow Sym$, ([1, Example 4.5]).

An important part of the structure of the Hopf algebra \mathcal{G} is the antipode $S: \mathcal{G} \rightarrow \mathcal{G}$. The cancellation-free formula for the antipode in terms of acyclic orientations of a graph Γ is obtained in [3]. We recall some basic definitions. Terminology comes from matroid theory. Given a graph $\Gamma = (V, E)$, for a collection of edges $F \subset E$ denote by $\Gamma_{V,F}$ the graph on V with the edge set F . A flat F of the graph Γ is a collection of its edges such that components of $\Gamma_{V,F}$ are induced subgraphs. The rank $\text{rk}(F)$ is the size of spanning forests of $\Gamma_{V,F}$. We have that $|V| = \text{rk}(F) + c(F)$, where $c(F)$ is the number of components of $\Gamma_{V,F}$. By contracting edges from a flat F we obtain the graph Γ/F . Finally, let $a(\Gamma)$ be the number of acyclic orientations of Γ . The formula of Humpert and Martin is as follows

$$S(\Gamma) = \sum_{F \in \mathcal{F}(\Gamma)} (-1)^{c(F)} a(\Gamma/F) \Gamma_{V,F}, \tag{3.1}$$

where the sum is over the set of flats $\mathcal{F}(\Gamma)$.

The following modification of the character ζ is considered in [2] in a wider context of the combinatorial Hopf algebra of simplicial complexes. Define $\zeta_q(\Gamma) = q^{\text{rk}(\Gamma)}$, which determines the algebra morphism $\zeta_q: \mathcal{G} \rightarrow \mathbf{k}(q)$, where $\mathbf{k}(q)$ is the field of rational functions in q . This character produces the unique morphism $\Psi_q: \mathcal{G} \rightarrow QSym$ to quasisymmetric functions over $\mathbf{k}(q)$. The expansion of $\Psi_q(\Gamma)$ in the monomial basis of quasisymmetric functions is determined by the universal formula [1, Theorem 4.1]

$$\Psi_q(\Gamma) = \sum_{\alpha \models n} (\zeta_q)_\alpha(\Gamma) M_\alpha.$$

The sum above is over all compositions of the integer $n = |V|$ and the coefficient of the expansion corresponding to the composition $\alpha = (a_1, \dots, a_k) \models n$ is given by

$$(\zeta_q)_\alpha(\Gamma) = \sum_{I_1 \sqcup \dots \sqcup I_k = V} q^{\text{rk}(\Gamma|_{I_1}) + \dots + \text{rk}(\Gamma|_{I_k})},$$

where the sum is over all set compositions of V of the type α . The coefficients $(\zeta_q)_\alpha(\Gamma)$ depend only on the partition corresponding to a composition α , so the function $\Psi_q(\Gamma)$ is actually symmetric and it can be expressed in the monomial basis of symmetric functions.

The invariant $\Psi_q(\Gamma)$ is more subtle than $\Psi(\Gamma)$. Obviously $\Psi_0(\Gamma)$ is the chromatic symmetric function of a graph Γ . It remains open to find two nonisomorphic graphs Γ_1 and Γ_2 with the same q -chromatic symmetric functions $\Psi_q(\Gamma_1) = \Psi_q(\Gamma_2)$. Let

$$\chi_q(\Gamma, d) = \text{ps}(\Psi_q(\Gamma))(d)$$

be the q -analog of the chromatic polynomial $\chi(\Gamma, d)$. It is a consequence of a general fact for combinatorial Hopf algebras (see [1]) that

$$\chi_q(\Gamma, -1) = (\zeta_q \circ S)(\Gamma). \tag{3.2}$$

Example 3.1. Consider the graph Γ on four vertices with the edge set $E = \{12, 13, 23, 34\}$. We find that

$$\Psi_q(\Gamma) = 24m_{1,1,1,1} + (8q + 4)m_{2,1,1} + (2q^2 + 4q)m_{2,2} + (3q^2 + q)m_{3,1} + q^3m_4.$$

By principal specialization and taking into account that

$$\text{ps}(m_{\lambda_{i_1}^{i_1}, \dots, \lambda_{i_k}^{i_k}})(d) = \frac{(i_1 + \dots + i_k)!}{i_1! \dots i_k!} \binom{d}{i_1 + \dots + i_k},$$

we obtain

$$\chi_q(\Gamma, d) = d(d-1)^2(d-2) + qd(d-1)(4d-5) + 4q^2d(d-1) + q^3d,$$

which by Theorem 1.1 gives

$$f(\mathcal{Z}_\Gamma, q) = 12 + 18q + 8q^2 + q^3.$$

4 Proof of Theorem 1.1

Proof. By applying (3.2) and the formula for antipode (3.1) we obtain

$$(-1)^{|V|} \chi_{-q}(\Gamma, -1) = (-1)^{|V|} \sum_{F \in \mathcal{F}(\Gamma)} (-1)^{c(\Gamma)} a(\Gamma/F) (-q)^{\text{rk}(F)}.$$

It follows that the statement of the theorem is equivalent to the following expression of the f -polynomial

$$f(\mathcal{Z}_\Gamma, q) = \sum_{F \in \mathcal{F}(\Gamma)} a(\Gamma/F) q^{\text{rk}(F)}. \tag{4.1}$$

Therefore it should be shown that components of f -vectors are determined by

$$f_k(\mathcal{Z}_\Gamma) = \sum_{\substack{F \in \mathcal{F}(\Gamma) \\ \text{rk}(F)=k}} a(\Gamma/F), \quad 0 \leq k \leq n-1. \tag{4.2}$$

By duality between the face lattice of \mathcal{Z}_Γ and the face lattice of the fan $\mathcal{F}_{\mathcal{H}_\Gamma}$ we have

$$f_k(\mathcal{Z}_\Gamma) = f_{n-k-1}(\mathcal{F}_{\mathcal{H}_\Gamma}).$$

Let $L(\mathcal{H}_\Gamma)$ be the intersection lattice of the graphical arrangement \mathcal{H}_Γ . For a subspace $X \in L(\mathcal{H}_\Gamma)$ there is an arrangement of hyperplanes

$$\mathcal{H}_\Gamma^X = \{X \cap H \mid X \not\subseteq H, H \in \mathcal{H}_\Gamma\}$$

whose intersection lattice $L(\mathcal{H}_\Gamma^X)$ is isomorphic to the upper cone of X in $L(\mathcal{H}_\Gamma)$. Since \mathcal{H}_Γ is central and essential we have

$$f_{n-k-1}(\mathcal{F}_{\mathcal{H}_\Gamma}) = \sum_{\substack{X \in L(\mathcal{H}_\Gamma) \\ \dim(X)=n-k-1}} r(\mathcal{H}_\Gamma^X), \tag{4.3}$$

where $r(\mathcal{H}_\Gamma^X)$ is the number of regions of the arrangement \mathcal{H}_Γ^X , see [8, Theorem 2.6].

The intersection lattice $L(\mathcal{H}_\Gamma)$ is isomorphic to the lattice of flats of the graphical matroid $M(\Gamma)$. By this isomorphism to a flat F of rank k corresponds the intersection subspace $X^F = \cap_{\{i,j\} \in F} H_{e_i - e_j}$ of dimension $n - k - 1$. It is easy to see that arrangements $\mathcal{H}_\Gamma^{X^F}$ and $\mathcal{H}_{\Gamma/F}$ coincide, which by (2.2) and comparing formulas (4.2) and (4.3) proves theorem. □

5 Examples

By applying Theorem 1.1 we obtain the following interpretation of identities elaborated in [2, Propositions 17, 19].

Example 5.1. (i) For the permutohedron $Pe^{n-1} = \mathcal{Z}_{K_n}$, the f -polynomial is given by

$$f(\mathcal{Z}_{K_n}, q) = A_n(q + 1),$$

where $A_n(q) = \sum_{\pi \in S_n} q^{\text{des}(\pi)}$ is the Euler polynomial. Recall that $\text{des}(\pi)$ is the number of descents of a permutation $\pi \in S_n$. It recovers the fact that the h -polynomial of the permutohedron Pe^{n-1} is the Euler polynomial $A_n(q)$.

(ii) For the cube $I^{n-1} = \mathcal{Z}_{T_n}$, where T_n is a tree on n vertices, the f -polynomial is given by

$$f(\mathcal{Z}_{T_n}, q) = (q + 2)^{n-1}.$$




Figure 2: Rhombic dodecahedron \mathcal{Z}_{C_4} .

Proposition 5.2. The f -polynomial of the graphical zonotope \mathcal{Z}_{C_n} associated to the cycle graph C_n on n vertices is given by

$$f(\mathcal{Z}_{C_n}, q) = q^n + q^{n-1} + (q + 2)^n - 2(q + 1)^n.$$

Proof. A flat $F \in \mathcal{F}(C_n)$ is determined by the complementary set of edges. If $\text{rk}(F) = n - k, k > 1$ then the complementary set has k edges and $C_n/F = C_k$. Since $a(C_k) = 2^k - 2, k > 1$, by formula (4.2), we obtain

$$f_{n-k}(\mathcal{Z}_{C_n}) = (2^k - 2) \binom{n}{k}, \quad 2 \leq k \leq n,$$

which leads to the required formula. □

Specially, for $n = 4$ the resulting zonotope is the rhombic dodecahedron (see Figure 2). We have

$$f(\mathcal{Z}_{C_4}, q) = 14 + 24q + 12q^2 + q^3.$$

Proposition 5.3. Let $\Gamma = \Gamma_1 \vee_v \Gamma_2$ be the wedge of two connected graphs Γ_1 and Γ_2 at the common vertex v . Then

$$f(\mathcal{Z}_\Gamma, q) = f(\mathcal{Z}_{\Gamma_1}, q)f(\mathcal{Z}_{\Gamma_2}, q).$$

Proof. The graphical matroids of involving graphs are related by $M(\Gamma) = M(\Gamma_1) \oplus M(\Gamma_2)$. For the sets of flats it holds $\mathcal{F}(\Gamma) = \{F_1 \cup F_2 \mid F_i \in \mathcal{F}(\Gamma_i), i = 1, 2\}$. For $F = F_1 \cup F_2$ we have $\Gamma/F = \Gamma_1/F_1 \vee_{[v]} \Gamma_2/F_2$, where $[v]$ is the component of the vertex v in $\Gamma_{V,F}$. Obviously $a(\Gamma/F) = a(\Gamma_1/F_1)a(\Gamma_2/F_2)$ and $\text{rk}(F) = \text{rk}(F_1) + \text{rk}(F_2)$. The proposition follows from formula (4.1). \square

The formula for cubes in Example 5.1 (ii) follows from Proposition 5.3 since any tree is a consecutive wedge of edges and $f(I^1, q) = q + 2$. It also allows us to restrict ourselves only to biconnected graphs. For a biconnected graph Γ with a disconnecting pair of vertices $\{u, v\}$ Whitney introduced the transformation called the *twist* around the pair $\{u, v\}$. This transformation does not have an affect on the graphical matroid $M(\Gamma)$ [9].




Figure 3: Biconnected graphs related by twist transformation.

Example 5.4. Figure 3 shows the pair of biconnected graphs on six vertices obtained one from another by the twist transformation. The corresponding zonotopes have the same f -polynomial

$$f(\mathcal{Z}_{\Gamma_1}, q) = f(\mathcal{Z}_{\Gamma_2}, q) = 126 + 348q + 358q^2 + 164q^3 + 30q^4 + q^5.$$

On the other hand their q -chromatic symmetric functions are different. One can check that corresponding coefficients by $m_{3,1^3}$ are different

$$\begin{aligned} [m_{3,1^3}] \Psi_q(\Gamma_1) &= (11q^2 + 8q + 1) \cdot 3!, \\ [m_{3,1^3}] \Psi_q(\Gamma_2) &= (10q^2 + 10q) \cdot 3!. \end{aligned}$$

This shows that the q -analog of the chromatic symmetric function of a graph is not determined by the corresponding graphical matroid. By taking $q = 0$ we obtain that even the chromatic symmetric functions are different since $[m_{3,1^3}] \Psi(\Gamma_1) = 6$ and $[m_{3,1^3}] \Psi(\Gamma_2) = 0$.

Let us now consider Stanley’s example of nonisomorphic graphs with the same chromatic symmetric functions, see [7]. We find that the f -polynomials of the corresponding graphical zonotopes differ for those graphs. From these examples we conclude that chromatic properties of a graph and the f -vector of the corresponding graphical zonotope are not related.

We have already noted that graphical zonotopes are generalized permutohedra. The h -polynomials of simple generalized permutohedra are determined in [5, Theorem 4.2]. The only simple graphical zonotopes are products of permutohedra [5, Proposition 5.2]. They are characterized by graphs whose biconnected components are complete subgraphs. Therefore Proposition 5.3 together with Example 5.1 (i) prove that the h -polynomial of a

simple graphical zonotope is the product of Eulerian polynomials, the fact obtained in [5, Corollary 5.4]. Example 3.1 is of this sort and represents the hexagonal prism which is the product $\mathcal{Z}_{K_3} \times \mathcal{Z}_{K_2}$.

References

- [1] M. Aguiar, N. Bergeron and F. Sottile, Combinatorial Hopf algebras and generalized Dehn-Sommerville relations, *Compos. Math.* **142** (2006), 1–30, doi:10.1112/s0010437x0500165x.
- [2] C. Benedetti, J. Hallam and J. Machacek, Combinatorial Hopf algebras of simplicial complexes, *SIAM J. Discrete Math.* **30** (2016), 1737–1757, doi:10.1137/15m1038281.
- [3] B. Humpert and J. L. Martin, The incidence Hopf algebra of graphs, *SIAM J. Discrete Math.* **26** (2012), 555–570, doi:10.1137/110820075.
- [4] A. Postnikov, Permutohedra, associahedra, and beyond, *Int. Math. Res. Not.* **2009** (2009), 1026–1106, doi:10.1093/imrn/rnn153.
- [5] A. Postnikov, V. Reiner and L. Williams, Faces of generalized permutohedra, *Doc. Math.* **13** (2008), 207–273, <http://www.math.uiuc.edu/documenta/vol-13/10.html>.
- [6] R. P. Stanley, Acyclic orientations of graphs, *Discrete Math.* **5** (1973), 171–178, doi:10.1016/0012-365x(73)90108-8.
- [7] R. P. Stanley, A symmetric function generalization of the chromatic polynomial of a graph, *Adv. Math.* **111** (1995), 166–194, doi:10.1006/aima.1995.1020.
- [8] R. P. Stanley, An introduction to hyperplane arrangements, in: E. Miller, V. Reiner and B. Sturmfels (eds.), *Geometric Combinatorics*, American Mathematical Society, Providence, Rhode Island, volume 13 of *IAS/Park City Mathematics Series*, pp. 389–496, 2007.
- [9] H. Whitney, 2-isomorphic graphs, *Amer. J. Math.* **55** (1933), 245–254, doi:10.2307/2371127.
- [10] G. M. Ziegler, *Lectures on Polytopes*, volume 152 of *Graduate Texts in Mathematics*, Springer-Verlag, New York, 1995, doi:10.1007/978-1-4613-8431-1.