

# The 4-girth-thickness of the complete graph

Christian Rubio-Montiel

*UMI LAFMIA 3175 CNRS at CINVESTAV-IPN, 07300, Mexico City, Mexico  
División de Matemáticas e Ingeniería, FES Ac-UNAM, 53150, Naulcalpan, Mexico*

Received 10 March 2017, accepted 21 June 2017, published online 19 September 2017

---

## Abstract

In this paper, we define the 4-girth-thickness  $\theta(4, G)$  of a graph  $G$  as the minimum number of planar subgraphs of girth at least 4 whose union is  $G$ . We prove that the 4-girth-thickness of an arbitrary complete graph  $K_n$ ,  $\theta(4, K_n)$ , is  $\lceil \frac{n+2}{4} \rceil$  for  $n \neq 6, 10$  and  $\theta(4, K_6) = 3$ .

*Keywords:* Thickness, planar decomposition, girth, complete graph.

*Math. Subj. Class.:* 05C10

---

## 1 Introduction

A finite graph  $G$  is *planar* if it can be embedded in the plane without any two of its edges crossing. A planar graph of order  $n$  and girth  $g$  has size at most  $\frac{g}{g-2}(n-2)$  (see [6]), and an acyclic graph of order  $n$  has size at most  $n-1$ , in this case, we define its girth as  $\infty$ . The *thickness*  $\theta(G)$  of a graph  $G$  is the minimum number of planar subgraphs whose union is  $G$ ; i.e. the minimum number of planar subgraphs into which the edges of  $G$  can be partitioned.

The thickness was introduced by Tutte [11] in 1963. Since then, exact results have been obtained when  $G$  is a complete graph [1, 3, 4], a complete multipartite graph [5, 12, 13] or a hypercube [9]. Also, some generalizations of the thickness for the complete graph  $K_n$  have been studied such that the outerthickness  $\theta_o$ , defined similarly but with outerplanar instead of planar [8], and the  $S$ -thickness  $\theta_S$ , considering the thickness on a surfaces  $S$  instead of the plane [2]. See also the survey [10].

We define the  $g$ -girth-thickness  $\theta(g, G)$  of a graph  $G$  as the minimum number of planar subgraphs of girth at least  $g$  whose union is  $G$ . Note that the 3-girth-thickness  $\theta(3, G)$  is the usual thickness and the  $\infty$ -girth-thickness  $\theta(\infty, G)$  is the *arboricity number*, i.e. the minimum number of acyclic subgraphs into which  $E(G)$  can be partitioned. In this paper, we obtain the 4-girth-thickness of an arbitrary complete graph of order  $n \neq 10$ .

---

*E-mail address:* christian@cs.cinvestav.mx (Christian Rubio-Montiel)

## 2 The exact value of $\theta(4, K_n)$ for $n \neq 10$

Since the complete graph  $K_n$  has size  $\binom{n}{2}$  and a planar graph of order  $n$  and girth at least 4 has size at most  $2(n - 2)$  for  $n \geq 3$  and  $n - 1$  for  $n \in \{1, 2\}$  then the 4-girth-thickness of  $K_n$  is at least

$$\left\lceil \frac{n(n-1)}{2(2n-4)} \right\rceil = \left\lceil \frac{n+1}{4} + \frac{1}{2n-4} \right\rceil = \left\lceil \frac{n+2}{4} \right\rceil$$

for  $n \geq 3$  and also  $\lceil \frac{n+2}{4} \rceil$  for  $n \in \{1, 2\}$ , we have the following theorem.

**Theorem 2.1.** *The 4-girth-thickness  $\theta(4, K_n)$  of  $K_n$  equals  $\lceil \frac{n+2}{4} \rceil$  for  $n \neq 6, 10$  and  $\theta(4, K_6) = 3$ .*

*Proof.* Figure 1 displays equality for  $n \leq 5$ .




Figure 1:  $\theta(4, K_n) = \lceil \frac{n+2}{4} \rceil$  for  $n = 1, 2, 3, 4, 5$ .

To prove that  $\theta(4, K_6) = 3 > \lceil \frac{6+2}{4} \rceil = 2$ , suppose that  $\theta(4, K_6) = 2$ . This partition define an edge coloring of  $K_6$  with two colors. By Ramsey’s Theorem, some part contains a triangle obtaining a contradiction for the girth 4. Figure 2 shows a partition of  $K_6$  into tree planar subgraphs of girth at least 4.




Figure 2:  $\theta(4, K_6) = 3$ .

For the remainder of this proof, we need to distinguish four cases, namely, when  $n = 4k - 1, n = 4k, n = 4k + 1$  and  $n = 4k + 2$  for  $k \geq 2$ . Note that in each case, the lower bound of the 4-girth thickness require at least  $k + 1$  elements. To prove our theorem, we exhibit a decomposition of  $K_{4k}$  into  $k + 1$  planar graphs of girth at least 4. The other three

cases are based in this decomposition. The case of  $n = 4k - 1$  follows because  $K_{4k-1}$  is a subgraph of  $K_{4k}$ . For the case of  $n = 4k + 2$ , we add two vertices and some edges to the decomposition obtained in the case of  $n = 4k$ . The last case follows because  $K_{4k+1}$  is a subgraph of  $K_{4k+2}$ . In the proof, all sums are taken modulo  $2k$ .

1. Case  $n = 4k$ . It is well-known that a complete graph of even order contains a cyclic factorization of Hamiltonian paths, see [7]. Let  $G$  be a subgraph of  $K_{4k}$  isomorphic to  $K_{2k}$ . Label its vertex set  $V(G)$  as  $\{v_1, v_2, \dots, v_{2k}\}$ . Let  $\mathcal{F}_1$  be the Hamiltonian path with edges

$$v_1 v_2, v_2 v_{2k}, v_{2k} v_3, v_3 v_{2k-1}, \dots, v_{2+k} v_{1+k}.$$

Let  $\mathcal{F}_i$  be the Hamiltonian path with edges

$$v_i v_{i+1}, v_{i+1} v_{i-1}, v_{i-1} v_{i+2}, v_{i+2} v_{i-2}, \dots, v_{i+k+1} v_{i+k},$$

where  $i \in \{2, 3, \dots, k\}$ .

Such factorization of  $G$  is the partition  $\{E(\mathcal{F}_1), E(\mathcal{F}_2), \dots, E(\mathcal{F}_k)\}$ . We remark that the center of  $\mathcal{F}_i$  has the edge  $e = v_{i+\lceil \frac{k}{2} \rceil} v_{i+\lceil \frac{3k}{2} \rceil}$ , see Figure 3.




Figure 3: The Hamiltonian path  $\mathcal{F}_i$ : Left a): The dashed edge  $e$  for  $k$  odd. Right b) The dashed edge  $e$  for  $k$  even.

Now, consider the complete subgraph  $G'$  of  $K_{4k}$  such that  $G' = K_{4k} \setminus V(G)$ . Label its vertex set  $V(G')$  as  $\{v'_1, v'_2, \dots, v'_{2k}\}$  and consider the factorization, similarly as before,  $\{E(\mathcal{F}'_1), E(\mathcal{F}'_2), \dots, E(\mathcal{F}'_k)\}$  where  $\mathcal{F}'_i$  is the Hamiltonian path with edges

$$v'_i v'_{i+1}, v'_{i+1} v'_{i-1}, v'_{i-1} v'_{i+2}, v'_{i+2} v'_{i-2}, \dots, v'_{i+k+1} v'_{i+k},$$

where  $i \in \{1, 2, \dots, k\}$ .

Next, we construct the planar subgraphs  $G_1, G_2, \dots, G_{k-1}$  and  $G_k$  of girth 4, order  $4k$  and size  $8k - 4$  (observe that  $2(4k - 2) = 8k - 4$ ), and also the matching  $G_{k+1}$ , as follows. Let  $G_i$  be a spanning subgraph of  $K_{4k}$  with edges  $E(\mathcal{F}_i) \cup E(\mathcal{F}'_i)$  and

$$v_i v'_{i+1}, v'_i v_{i+1}, v_{i+1} v'_{i-1}, v'_{i+1} v_{i-1}, v_{i-1} v'_{i+2}, v'_{i-1} v_{i+2}, \dots, v_{i+k+1} v'_{i+k}, v'_{i+k+1} v_{i+k}$$

where  $i \in \{1, 2, \dots, k\}$ ; and let  $G_{k+1}$  be a perfect matching with edges  $v_j v'_j$  for  $j \in \{1, 2, \dots, 2k\}$ . Figure 4 shows  $G_i$  is a planar graph of girth at least 4.




Figure 4: Left a): The graph  $G_i$  for any  $i \in \{1, 2, \dots, k\}$ . Right b) The graph  $G_{k+1}$ .

To verify that  $K_{4k} = \bigcup_{i=1}^{k+1} G_i$ : 1) If the edge  $v_{i_1} v_{i_2}$  of  $G$  belongs to the factor  $\mathcal{F}_i$  then  $v_{i_1} v_{i_2}$  belongs to  $G_i$ . If the edge is primed, belongs to  $G'_i$ . 2) The edge  $v_{i_1} v'_{i_2}$  belongs to  $G_{k+1}$  if and only if  $i_1 = i_2$ , otherwise it belongs to the same graph  $G_i$  as  $v_{i_1} v_{i_2}$ . Similarly in the case of  $v'_{i_1} v_{i_2}$  and the result follows.

2. Case  $n = 4k - 1$ . Since  $K_{4k-1} \subset K_{4k}$ , we have

$$k + 1 \leq \theta(4, K_{4k-1}) \leq \theta(4, K_{4k}) \leq k + 1.$$

3. Case  $n = 4k + 2$  (for  $k \neq 2$ ). Let  $\{G_1, \dots, G_{k+1}\}$  be the planar decomposition of  $K_{4k}$  constructed in the Case 1. We will add the two new vertices  $x$  and  $y$  to every planar subgraph  $G_i$ , when  $1 \leq i \leq k + 1$ , and we will add 4 edges to each  $G_i$ , when  $1 \leq i \leq k$ , and  $4k + 1$  edges to  $G_{k+1}$  such that the resulting new subgraphs of  $K_{4k+2}$  will be planar. Note that  $\binom{4k}{2} + 4k + 4k + 1 = \binom{4k+2}{2}$ .

To begin with, we define the graph  $H_{k+1}$  adding the vertices  $x$  and  $y$  to the planar subgraph  $G_{k+1}$  and the  $4k + 1$  edges

$$\{xy, xv_1, xv'_2, xv_3, xv'_4, \dots, xv_{2k-1}, xv'_{2k}, yv'_1, yv_2, yv'_3, yv_4, \dots, yv'_{2k-1}, yv_{2k}\}.$$

The graph  $H_{k+1}$  has girth 4, see Figure 5.

In the following, for  $1 \leq i \leq k$ , by adding vertices  $x$  and  $y$  to  $G_i$  and adding 4 edges to  $G_i$ , we will get a new planar graph  $H_i$  such that  $\{H_1, \dots, H_{k+1}\}$  is a planar decomposition of  $K_{4k+2}$  such that the girth of every element is 4. To achieve it, the given edges to the graph  $H_i$  will be  $v'_j x, xv_{j-1}, v_j y, yv'_{j-1}$ , for some odd  $j \in \{1, 3, \dots, 2k - 1\}$ .

According to the parity of  $k$ , we have two cases:




Figure 5: The graph  $H_{k+1}$ .

- Suppose  $k$  odd. For odd  $i \in \{1, 2, \dots, k\}$ , we define the graph  $H_i$  adding the vertices  $x$  and  $y$  to the planar subgraph  $G_i$  and the 4 edges

$$\{xv'_{i+\lceil \frac{3k}{2} \rceil - 1}, xv_{i+\lceil \frac{3k}{2} \rceil}, yv_{i+\lceil \frac{3k}{2} \rceil - 1}, yv'_{i+\lceil \frac{3k}{2} \rceil}\}$$

when  $\lceil \frac{k}{2} \rceil$  is even, otherwise

$$\{yv'_{i+\lceil \frac{3k}{2} \rceil - 1}, yv_{i+\lceil \frac{3k}{2} \rceil}, xv_{i+\lceil \frac{3k}{2} \rceil - 1}, xv'_{i+\lceil \frac{3k}{2} \rceil}\}.$$

Additionally, for even  $i \in \{1, 2, \dots, k\}$ , we define the graph  $H_i$  adding the vertices  $x$  and  $y$  to the planar subgraph  $G_i$  and the 4 edges

$$\{xv'_{i+\lceil \frac{k}{2} \rceil - 1}, xv_{i+\lceil \frac{k}{2} \rceil}, yv_{i+\lceil \frac{k}{2} \rceil - 1}, yv'_{i+\lceil \frac{k}{2} \rceil}\}$$

when  $\lceil \frac{k}{2} \rceil$  is even, otherwise

$$\{yv'_{i+\lceil \frac{k}{2} \rceil - 1}, yv_{i+\lceil \frac{k}{2} \rceil}, xv_{i+\lceil \frac{k}{2} \rceil - 1}, xv'_{i+\lceil \frac{k}{2} \rceil}\}.$$

Note that the graph  $H_i$  has girth 4 for all  $i$ , see Figure 6.

- Suppose  $k$  even. Similarly that the previous case, for odd  $i \in \{1, 2, \dots, k\}$ , we define the graph  $H_i$  adding the vertices  $x$  and  $y$  to the planar subgraph  $G_i$  and the 4 edges

$$\{xv_{i+\lceil \frac{3k}{2} \rceil + 1}, xv'_{i+\lceil \frac{3k}{2} \rceil}, yv'_{i+\lceil \frac{3k}{2} \rceil + 1}, yv_{i+\lceil \frac{3k}{2} \rceil}\}$$

when  $\lceil \frac{k}{2} \rceil$  is even, otherwise

$$\{yv_{i+\lceil \frac{3k}{2} \rceil + 1}, yv'_{i+\lceil \frac{3k}{2} \rceil}, xv'_{i+\lceil \frac{3k}{2} \rceil + 1}, xv_{i+\lceil \frac{3k}{2} \rceil}\}.$$

On the other hand, for even  $i \in \{1, 2, \dots, k\}$ , we define the graph  $H_i$  adding the vertices  $x$  and  $y$  to the planar subgraph  $G_i$  and the 4 edges

$$\{xv_{i+\lceil \frac{k}{2} \rceil}, xv'_{i+\lceil \frac{k}{2} \rceil - 1}, yv'_{i+\lceil \frac{k}{2} \rceil}, yv_{i+\lceil \frac{k}{2} \rceil - 1}\}$$

when  $\lceil \frac{k}{2} \rceil$  is even, otherwise

$$\{yv_{i+\lceil \frac{k}{2} \rceil}, yv'_{i+\lceil \frac{k}{2} \rceil - 1}, xv'_{i+\lceil \frac{k}{2} \rceil}, xv_{i+\lceil \frac{k}{2} \rceil - 1}\}.$$

Note that the graph  $H_i$  has girth 4 for all  $i$ , see Figure 7.




Figure 6: The graph  $H_i$  when  $k$  is odd and its auxiliary graph  $\mathcal{F}_i^*$ . Above a) When  $i$  is odd. Bottom b) When  $i$  is even.




Figure 7: The graph  $H_i$  when  $k$  is even and its auxiliary graph  $\mathcal{F}_i^*$ . Above a) When  $i$  is odd. Bottom b) When  $i$  is even.

In order to verify that each edge of the set

$$\{xv'_1, xv_2, xv'_3, xv_3, \dots, xv'_{2k-1}, xv_{2k}, yv_1, yv'_2, yv_3, yv'_3, \dots, yv_{2k-1}, yv'_{2k}\}.$$

is in exactly one subgraph  $H_i$ , for  $i \in \{1, \dots, k\}$ , we obtain the unicyclic graph  $\mathcal{F}_i^*$  identifying  $v_j$  and  $v'_j$  resulting in  $v_j$ ; identifying  $x$  and  $y$  resulting in a vertex which is contracted with one of its neighbours. The resulting edge, in dashed, is showed in Figures 6 and 7. The set of those edges are a perfect matching of  $K_{2k}$  proving that the added two paths of length 2 in  $G_i$  have end vertices  $v_j$  and  $v'_{j-1}$ , and the other  $v'_j$  and  $v_{j-1}$ . The election of the label of the center vertex is such that one path is  $v_{even}xv'_{odd}$  and  $v'_{even}yv_{odd}$  and the result follows.

4. Case  $n = 4k + 1$  (for  $k \neq 2$ ). Since  $K_{4k+1} \subset K_{4k+2}$ , we have

$$k + 1 \leq \theta(4, K_{4k+1}) \leq \theta(4, K_{4k+2}) \leq k + 1.$$

For  $k = 2$ , Figure 8 displays a decomposition of three planar graphs of girth at least 4 proving that  $\theta(4, K_9) = \lceil \frac{9+2}{4} \rceil = 3$ .




Figure 8: A planar decomposition of  $K_9$  into three subgraphs of girth 4 and 5.

By the four cases, the theorem follows. □

About the case of  $K_{10}$ , it follows  $3 \leq \theta(4, K_{10}) \leq 4$ . We conjecture that  $\theta(4, K_{10}) = 4$ .

### References

- [1] V. B. Alekseev and V. S. Gončakov, The thickness of an arbitrary complete graph, *Mat. Sb. (N.S.)* **101(143)** (1976), 212–230.
- [2] L. W. Beineke, Minimal decompositions of complete graphs into subgraphs with embeddability properties, *Canad. J. Math.* **21** (1969), 992–1000.
- [3] L. W. Beineke and F. Harary, On the thickness of the complete graph, *Bull. Amer. Math. Soc.* **70** (1964), 618–620.
- [4] L. W. Beineke and F. Harary, The thickness of the complete graph, *Canad. J. Math.* **17** (1965), 850–859.
- [5] L. W. Beineke, F. Harary and J. W. Moon, On the thickness of the complete bipartite graph, *Proc. Cambridge Philos. Soc.* **60** (1964), 1–5.
- [6] J. Bondy and U. Murty, *Graph theory*, volume 244 of *Graduate Texts in Mathematics*, Springer, New York, 2008.



- [7] G. Chartrand and P. Zhang, *Chromatic graph theory*, Discrete Mathematics and its Applications, CRC Press, 2009.
- [8] R. K. Guy and R. J. Nowakowski, The outerthickness & outercoarseness of graphs. I. The complete graph & the  $n$ -cube, in: *Topics in combinatorics and graph theory (Oberwolfach, 1990)*, Physica, Heidelberg, pp. 297–310, 1990.
- [9] M. Kleinert, Die Dicke des  $n$ -dimensionalen Würfel-Graphen, *J. Combin. Theory* **3** (1967), 10–15.
- [10] P. Mutzel, T. Odenthal and M. Scharbrodt, The thickness of graphs: a survey, *Graphs Combin.* **14** (1998), 59–73.
- [11] W. T. Tutte, The thickness of a graph, *Indag. Math.* **25** (1963), 567–577.
- [12] Y. Yang, A note on the thickness of  $K_{l,m,n}$ , *Ars Combin.* **117** (2014), 349–351.
- [13] Y. Yang, Remarks on the thickness of  $K_{n,n,n}$ , *Ars Math. Contemp.* **12** (2017), 135–144.