

Arc-transitive graphs of valency 8 have a semiregular automorphism*

Gabriel Verret[†]

Centre for Mathematics of Symmetry and Computation, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

Faculty of Mathematics, Natural Sciences and Information Technologies, University of Primorska, Glagoljaška 8, 6000 Koper, Slovenia.

Received 23 May 2013, accepted 16 October 2013, published online 14 April 2014

Abstract

One version of the polycirculant conjecture states that every vertex-transitive graph has a non-identity semiregular automorphism that is, a non-identity automorphism whose cycles all have the same length. We give a proof of the conjecture in the arc-transitive case for graphs of valency 8, which was the smallest open valency.

Keywords: Arc-transitive graphs, polycirculant conjecture, semiregular automorphism.

Math. Subj. Class.: 20B25, 05E18

1 Introduction

All graphs considered in this paper are finite and simple. An automorphism of a graph Γ is a permutation of the vertex-set $V(\Gamma)$ which preserves the adjacency relation. The set of automorphisms of Γ forms a group, denoted $\text{Aut}(\Gamma)$. A graph Γ is said to be G -vertex-transitive if G is a subgroup of $\text{Aut}(\Gamma)$ acting transitively on $V(\Gamma)$. Similarly, Γ is said to be G -arc-transitive if G acts transitively on the arcs of Γ . (An *arc* is an ordered pair of adjacent vertices). When $G = \text{Aut}(\Gamma)$, the prefix G in the above notation is sometimes omitted.

A group G acting on a set Ω is *semiregular* on Ω if the stabiliser G_ω is trivial for every $\omega \in \Omega$; note that this implies that the action is faithful. An element $g \in G$ is called *semiregular* provided that it generates a semiregular group. Equivalently, an element $g \in G$ is semiregular if $\langle g \rangle$ acts faithfully on Ω and that all of the cycles of the permutation induced by g have the same length.

*Dedicated to Dragan Marušič on the occasion of his 60th birthday.

[†]The author is supported by UWA as part of the Australian Research Council grant DE130101001.

E-mail address: gabriel.verret@uwa.edu.au (Gabriel Verret)

In 1981, Marušič conjectured [7] that every vertex-transitive graph with at least two vertices has a non-identity semiregular automorphism. This is sometimes called the *polycirculant conjecture*. While there has been a lot of work on this conjecture and some of its variants, it is still wide open. There has been progress in some directions (see [6] for a survey). For example, the conjecture has been settled for graphs of certain orders [2, 7, 8] and for graphs of valency at most four [1, 8].

In the arc-transitive case, slightly more can be said, but we first need a few definitions. If Γ is a graph and $G \leq \text{Aut}(\Gamma)$ then $G_v^{\Gamma(v)}$ denotes the permutation group induced by the action of the vertex-stabiliser G_v on the neighbourhood $\Gamma(v)$ of the vertex v . A permutation group is called *quasiprimitive* if each of its nontrivial normal subgroup is transitive, while a vertex-transitive graph Γ is called *locally-quasiprimitive* if $\text{Aut}(\Gamma)_v^{\Gamma(v)}$ is quasiprimitive. Using this terminology, we have the following theorem due to Giudici and Xu.

Theorem 1.1. [5, Theorem 1.1] *A locally-quasiprimitive vertex-transitive graph has a non-identity semiregular automorphism.*

A transitive group of prime degree is clearly quasiprimitive (in fact primitive) and hence it follows from Theorem 1.1 that arc-transitive graphs of prime valency have a non-identity semiregular automorphism. In an upcoming paper [4], the author and Giudici deal with the case of arc-transitive graphs of valency twice a prime. The main result of this paper is to prove the polycirculant conjecture for arc-transitive graphs of valency 8, the smallest open valency.

Theorem 1.2. *An arc-transitive graph of valency 8 has a non-identity semiregular automorphism.*

The prime 2 plays a special role in our proof. In some sense, we use the fact that any proper factorisation of 8 must include 2 as one of the factors. In particular, if Γ is a G -arc-transitive graph of valency 8 and N is a normal subgroup of G that is not semiregular and that has at least three orbits then the quotient graph Γ/N is an arc-transitive graph of valency either 2 or 4. In the former case, we can view Γ/N as an asymmetric arc-transitive digraph of out-valency 4, which turns out to be very useful (see Theorem 2.7). In the latter case, the valency of Γ is exactly twice the valency of Γ/N which gives us very precious information about the kernel of the action of G on N -orbits (see the proof of Theorem 1.2).

This special role of the prime 2 is also evident in the proof of the case of valency twice a prime [4]. In the more general case of valency a product of two primes, Xu [13] has made significant progress and it seems the bottleneck is the case when G is solvable. (See [3] for some results in this case.)

2 Preliminaries

2.1 Permutation groups

We start with a few very basic lemmas about permutation groups. Since the proofs are short, we include them for the sake of completeness.

Lemma 2.1. *Let p be a prime and let G be a transitive permutation group of degree a power of p . Then G contains a non-identity semiregular element.*

Proof. Let S be a Sylow p -subgroup of G . By [12, Theorem 3.4'], S is transitive. Let z be a non-identity element of the centre of S . If z fixes a point then it must fix all of them, which is a contradiction. It follows that z is semiregular. \square

Lemma 2.2. *If G is a transitive permutation group with a non-trivial abelian normal subgroup N that has at most two orbits, then N contains a non-identity semiregular element.*

Proof. If N is semiregular then we are done. We may thus assume that there exists a non-identity element $n \in N$ fixing some point v . Since N is abelian, n fixes the N -orbit v^N pointwise. It follows that N has exactly two orbits. Let u be a point not in v^N and let $g \in G$ such that $v^g = u$. Then n acts semiregularly on u^N , while n^g fixes u^N pointwise and acts semiregularly on v^N . It follows that nn^g is a non-identity semiregular element of N . \square

Lemma 2.3. *Let G be a permutation group and let K be a normal subgroup of G such that G/K acts faithfully on the K -orbits. If G/K contains a semiregular element gK of order r coprime to $|K|$ then G contains a semiregular element of order r .*

Proof. There exists $k \in K$ such that $g^r = k$. Let $h = g^{|k|}$. Since $|k|$ and r are coprime, it follows that $|h| = r$ and hK is a non-identity power of gK hence it is semiregular. We now show that h is semiregular. Suppose h^i fixes a point v for some integer i . Then $h^i K = (hK)^i$ fixes v^K and, since hK is semiregular, $h^i K = K$. This implies that $h^i \in K$. Since h has order coprime to $|K|$, it follows that $h^i = 1$. \square

Lemma 2.4. *A transitive permutation group of degree 8 is either primitive or it is a $\{2, 3\}$ -group.*

Proof. Let G be a transitive permutation group of degree 8 that is not primitive. Then G has a non-trivial system of imprimitivity, with blocks of size either 2 or 4. It follows that G is isomorphic to a subgroup of $\text{Sym}(4) \wr \text{Sym}(2)$ or $\text{Sym}(2) \wr \text{Sym}(4)$ and hence it is a $\{2, 3\}$ -group. \square

We also need the following folklore lemma

Lemma 2.5. *Let p be a prime, let Γ be a connected graph and let $G \leq \text{Aut}(\Gamma)$. If, for every $v \in V(\Gamma)$, $|G_v^{\Gamma(v)}|$ is coprime to p , then so is $|G_u|$ for every $u \in V(\Gamma)$.*

Proof. We prove this by contradiction. Let p be a prime dividing $|G_u|$ for some $u \in V(\Gamma)$ and let $g \in G_u$ have order p . Since g is non-trivial, it must move some vertex. Let w be a vertex of Γ moved by g at minimal distance from u . By the connectivity of Γ , there is a path u, u_1, \dots, u_t, w such that g fixes each u_i . Then $g \in G_{u_t}$ and g acts nontrivially on $\Gamma(u_t)$. Thus p divides $|G_{u_t}^{\Gamma(u_t)}|$ and the result follows. \square

2.2 Quotient graphs

We will need some basic facts about quotient graphs, which we now collect. (For a reference, see [11] for example.)

Let Γ be a graph and let $N \leq \text{Aut}(\Gamma)$. The *quotient graph* Γ/N is the graph whose vertices are the N -orbits with two such N -orbits v^N and u^N adjacent whenever there is a pair of vertices $v' \in v^N$ and $u' \in u^N$ that are adjacent in Γ . Clearly, if Γ is connected then so is Γ/N .

Let Γ be a G -arc-transitive graph, let $N \trianglelefteq G$ and let K be the kernel of the action of G on N -orbits. Then $K = NK_v \trianglelefteq G$, $G/K \leq \text{Aut}(\Gamma/N)$ and Γ/N is G/K -arc-transitive. If N has at least 3 orbits then the valency of Γ/N is at least 2 and divides the valency of Γ . If Γ/N has the same valency as Γ then $N = K$ and K is semiregular.

Some of these facts are used in the proof of the following lemma.

Lemma 2.6. *Let p be an odd prime and let Γ be a connected 4-valent G -arc-transitive graph such that p divides $|\mathbb{V}(\Gamma)|$. If G is solvable then G contains a semiregular element of order p .*

Proof. The proof goes by induction on $|\mathbb{V}(\Gamma)|$. Let $v \in \mathbb{V}(\Gamma)$. If $G_v^{\Gamma(v)}$ is a 2-group then it follows from Lemma 2.5 that G_v is a 2-group and every element of G of order p is semiregular. We thus assume that $G_v^{\Gamma(v)}$ is not a 2-group and, since it is a transitive group of degree 4, it is 2-transitive. Since G is solvable, it contains a non-trivial normal elementary abelian q -group N . If N has at most two orbits then $p = q$ and the result follows from Lemma 2.2.

From now on, we assume that N has at least three orbits. Since $G_v^{\Gamma(v)}$ is 2-transitive, this implies that Γ/N is 4-valent and hence N is semiregular and G/N acts faithfully on Γ/N . If $p = q$ then N contains a semiregular element of order p and the conclusion holds. Suppose now that $p \neq q$. Then Γ/N is a connected, 4-valent G/N -arc-transitive graph. Since p is coprime to $|N|$, it divides $|\mathbb{V}(\Gamma/N)|$. By the induction hypothesis, G/N contains a semiregular element of order p . The result then follows from Lemma 2.3. \square

2.3 Digraphs

Finally, we will need a few notions about digraphs. We follow the terminology of [10] closely. A *digraph* $\vec{\Gamma}$ consists of a finite non-empty set of *vertices* $\mathbb{V}(\vec{\Gamma})$ and a set of *arcs* $A(\vec{\Gamma}) \subseteq V \times V$, which is an arbitrary binary relation on V . A digraph $\vec{\Gamma}$ is called *asymmetric* provided that the relation $A(\vec{\Gamma})$ is asymmetric.

If (u, v) is an arc of $\vec{\Gamma}$ then we say that v is an *out-neighbour* of u and that u is an *in-neighbour* of v . The symbols $\vec{\Gamma}^+(v)$ and $\vec{\Gamma}^-(v)$ will denote the set of out-neighbors of v and the set of in-neighbors of v , respectively. We also say that u is the *tail* and v the *head* of (u, v) , respectively. The digraph $\vec{\Gamma}$ is said to be of *out-valence* k if $|\vec{\Gamma}^+(v)| = k$ for every $v \in \mathbb{V}(\vec{\Gamma})$.

An automorphism of $\vec{\Gamma}$ is a permutation of $\mathbb{V}(\vec{\Gamma})$ which preserves the relation $A(\vec{\Gamma})$. The set of automorphisms of $\vec{\Gamma}$ forms a group, denoted $\text{Aut}(\vec{\Gamma})$. We say that $\vec{\Gamma}$ is *arc-transitive* provided that $\text{Aut}(\vec{\Gamma})$ acts transitively on $A(\vec{\Gamma})$.

We say that two arcs a and b of $\vec{\Gamma}$ are *related* if they have a common tail or a common head. Let R denote the transitive closure of this relation. The *alternet* of $\vec{\Gamma}$ (with respect to a) is the subdigraph of $\vec{\Gamma}$ induced by the R -equivalence class $R(a)$ of the arc a . (i.e. the digraph with vertex-set consisting of all heads and tails of arcs in $R(a)$ and whose arc-set is $R(a)$). If the alternet with respect to (u, v) contains an arc of the form (v, w) then this alternet is called *degenerate*.

If the alternet of the arc (u, v) is non-degenerate then it is a connected bipartite digraph where the first bipartition set consists only of sources while the second bipartition set contains only sinks. An important case occurs when this alternet is in fact a complete bipartite digraph in which case we will simply say that the alternet is *complete bipartite*. We say that $\vec{\Gamma}$ is *loosely attached* if $\vec{\Gamma}$ has no degenerate alternets and the intersection of the set of sinks of one alternet intersects the set of sources of another alternet in at most one vertex.

We define the *digraph of alternets* $\text{Al}(\vec{\Gamma})$ of $\vec{\Gamma}$ as the digraph the vertices of which are the alternets of $\vec{\Gamma}$ and with two alternets A and B forming an arc (A, B) of $\text{Al}(\vec{\Gamma})$ whenever the intersection of the set of sinks of A with the set of sources of B is non-empty.

We are now ready to prove the following theorem.

Theorem 2.7. *Let $\vec{\Gamma}$ be a connected asymmetric arc-transitive digraph of out-valence 4. Then $\vec{\Gamma}$ has a non-identity semiregular automorphism.*

Proof. The proof goes by induction on $|\mathcal{V}(\vec{\Gamma})|$. Since $\vec{\Gamma}$ is connected and arc-transitive (and finite), it follows that it is vertex-transitive and strongly connected (see for example [9, Lemma 2]).

Let $v \in \mathcal{V}(\vec{\Gamma})$ and let $G = \text{Aut}(\vec{\Gamma})$. Without loss of generality, we may assume that every prime that divides $|G|$ also divides $|G_v|$. If G_v is a 2-group then the conclusion follows from Lemma 2.1. We may thus assume that G_v is not a 2-group and hence neither is $G_v^{\Gamma^+(v)}$. Since $G_v^{\Gamma^+(v)}$ is a transitive permutation group of degree 4, it is 2-transitive.

Since $\vec{\Gamma}$ has out-valence 4, $G_v^{\Gamma^+(v)}$ is a $\{2, 3\}$ -group, hence so are G_v and G and therefore G is solvable by Burnside's Theorem. It follows that G has an abelian minimal normal subgroup N . If N is semiregular then the conclusion holds. We may thus assume that N is not semiregular and hence $N_v^{\Gamma^+(v)}$ is a non-trivial normal subgroup of $G_v^{\Gamma^+(v)}$ and therefore is transitive. The same argument yields that $N_v^{\Gamma^-(v)}$ is also transitive.

Let (u, v) be an arc of $\vec{\Gamma}$. We have just seen that $N_u^{\Gamma^+(u)}$ and $N_v^{\Gamma^-(v)}$ are both transitive. This implies that $\Gamma^+(u) \subseteq v^N$ and $\Gamma^-(v) \subseteq u^N$. On the other hand, N is abelian and hence N_u fixes u^N pointwise and N_v fixes v^N pointwise. It follows that the alternet of $\vec{\Gamma}$ with respect to (u, v) is not degenerate and is complete bipartite.

If $\vec{\Gamma}$ is not loosely attached then it follows that, for every vertex x , there exists at least one other vertex y such that $\vec{\Gamma}^-(x) = \vec{\Gamma}^-(y)$ and $\vec{\Gamma}^+(x) = \vec{\Gamma}^+(y)$. It follows easily that $\vec{\Gamma}$ has a non-identity semiregular automorphism in this case.

We may thus assume that $\vec{\Gamma}$ is loosely attached. Let $\vec{\Gamma}' = \text{Al}(\vec{\Gamma})$ be the digraph of alternet of $\vec{\Gamma}$. It follows from [10, Lemmas 3.1-3.3] that $\vec{\Gamma}'$ is a connected asymmetric digraph of out-valence 4 and that $G = \text{Aut}(\vec{\Gamma}')$. It also follows easily that an automorphism which is semiregular on $\vec{\Gamma}'$ is also semiregular on $\vec{\Gamma}$.

Note that $|\mathcal{V}(\vec{\Gamma}')| = |\mathcal{V}(\vec{\Gamma})|/4$ and hence we may apply the induction hypothesis to $\vec{\Gamma}'$ to conclude that it has a non-identity semiregular automorphism g . Together with the observation in the previous sentence, this concludes the proof. \square

3 Proof of Theorem 1.2

Let Γ be an arc-transitive graph of valency 8. We must show that Γ has a non-identity semiregular automorphism.

Clearly, we may assume that Γ is connected. Let $G = \text{Aut}(\Gamma)$. If Γ is locally-quasiprimitive then the result follows from Theorem 1.1. We therefore assume that $G_v^{\Gamma(v)}$ is not quasiprimitive. By Lemma 2.4, $G_v^{\Gamma(v)}$ is a $\{2, 3\}$ -group and hence, by Lemma 2.5, so is G_v . We may also assume that G itself is a $\{2, 3\}$ -group and hence it is solvable by Burnside's Theorem.

It follows that G has an elementary abelian minimal normal subgroup N . If N has at most two orbits then the result follows from Lemma 2.2. We may thus assume that N has at least three orbits and, in particular, $N_v^{\Gamma(v)}$ is intransitive. If N is semiregular then the conclusion follows. We may thus assume that $N_v \neq 1$ and hence $N_v^{\Gamma(v)} \neq 1$. In particular, Γ/N has valency strictly less than 8 and $N_v^{\Gamma(v)}$ is a non-trivial, intransitive normal subgroup

of $G_v^{\Gamma(v)}$. Since the orbits of $N_v^{\Gamma(v)}$ are blocks of $G_v^{\Gamma(v)}$ it follows that $N_v^{\Gamma(v)}$ is a 2-group and hence N is an elementary abelian 2-group. Let K be the kernel of the action of G on N -orbits.

If Γ/N is 4-valent then v is adjacent to at most two vertices from any N -orbit. It follows that $K_v^{\Gamma(v)}$ is a 2-group and hence so are K_v and K . If $|\mathbf{V}(\Gamma/N)|$ is a power of 2 then so is $|\mathbf{V}(\Gamma)|$ and the result follows from Lemma 2.1. We may thus assume that $|\mathbf{V}(\Gamma/N)|$ is not a power of 2 and, by Lemma 2.6, G/K contains a semiregular element of odd order. It then follows from Lemma 2.3 that G contains a semiregular element of odd order.

It remains to deal with the case when Γ/N is 2-valent. In this case, there is a natural orientation of Γ as a connected asymmetric 4-valent digraph $\vec{\Gamma}$ and $\text{Aut}(\vec{\Gamma})$ is a subgroup of index 2 in G . By Theorem 2.7, $\text{Aut}(\vec{\Gamma})$ has a semiregular element. This concludes the proof.

Acknowledgements. We thank the anonymous referees for their valuable advice.

References

- [1] E. Dobson, A. Malnič, D. Marušič and L. A. Nowitz, Semiregular automorphisms of vertex-transitive graphs of certain valencies, *J. Combin. Theory, Ser. B* **97** (2007), 371–380.
- [2] E. Dobson, A. Malnič, D. Marušič and L. A. Nowitz, Minimal normal subgroups of transitive permutation groups of square-free degree, *Discrete Math.* **307** (2007), 373–385.
- [3] E. Dobson and D. Marušič, On semiregular elements of solvable groups, *Comm. Algebra* **39** (2011), 1413–1426.
- [4] M. Giudici and G. Verret, Semiregular automorphisms in arc-transitive graphs of valency $2p$, in preparation.
- [5] M. Giudici and J. Xu, All vertex-transitive locally-quasiprimitive graphs have a semiregular automorphism, *J. Algebraic Combin.* **25** (2007), 217–232.
- [6] K. Kutnar and D. Marušič, Recent Trends and Future Directions in Vertex-Transitive Graphs, *Ars Math. Contemp.* **1** (2008), 112–125.
- [7] D. Marušič, On vertex symmetric digraphs, *Discrete Math.* **36** (1981), 69–81.
- [8] D. Marušič and R. Scapellato, Permutation groups, vertex-transitive digraphs and semiregular automorphisms, *European J. Combin.* **19** (1998), 707–712.
- [9] P. M. Neumann, Finite Permutation Groups, edge-coloured graphs and matrices, in: M. P. J. Curran (ed.), *Topics in groups theory and computations*, Academic Press, London (1977).
- [10] P. Potočnik and G. Verret, On the vertex-stabiliser in arc-transitive digraphs, *J. Combin. Theory, Ser. B* **100** (2010), 497–509.
- [11] C. E. Praeger, Imprimitive symmetric graphs, *Ars Combin.* **19** (1985), 149–163.
- [12] H. Wielandt, *Finite permutation groups*, translated from German by R. Bercov, Academic Press, New York (1964).
- [13] J. Xu, Semiregular automorphisms of arc-transitive graphs with valency pq , *European J. Combin.* **29** (2008), 622–629.