Complete graphs with zero diagonal inverse

Alexander Farrugia , John Baptist Gauci , Irene Sciriha *

Department of Mathematics, Faculty of Science, University of Malta

Received 19 March 2014, accepted 23 October 2014, published online 15 November 2015

Abstract

If the inverse of a non-singular real symmetric matrix that represents an edge-weighted graph with no loops has zero diagonal, then the inverse itself is the matrix of a loopless graph. Here we show that such graphs having non-zero weight on each edge always exist if their number of vertices is at least 6.

Keywords: NSSD, $\mathbf{G}$ -nutful graph, circulant matrix, complete graph.

Math. Subj. Class.: 05C90, 05B20

1 Introduction

Since the discovery of fullerenes in 1985, an intimate relationship has matured between these carbon molecules and mathematics. The combinatorial, graph theoretical and algebraic properties of fullerenes have been exploited by many researchers (see, for example [1, 3]) to classify the structures of fullerenes and to predict the unique chemical and physical properties of these highly symmetric structures. In “classical” fullerenes, the structure is composed of solely pentagonal and hexagonal regions joined together to form 3-regular polyhedra [3]. However, over the years, non-traditional fullerenes composed of cycles of other sizes, which are general 3-regular polyhedra, have been studied, especially for their regularity properties. This current work is motivated mainly by the beauty of these structures, coupled with the elegance of the interplay between combinatorics, graph theory and algebra. In particular, we consider the algebraic and structural properties of a special family of graphs, termed $\mathbf{G}$ -nutful graphs, introduced in [8].

The graphs we consider have weighted edges and no loops. If a non-singular real symmetric matrix $\mathbf{G}$ representing such a graph has an inverse $\mathbf{G}^{-1}$ with each entry of its diagonal equal to zero, then the graph is referred to as a NSSD (*Non-Singular graphs with*

*Corresponding author.

E-mail addresses: alex.farrugia@um.edu.mt (Alexander Farrugia), john-baptist.gauci@um.edu.mt (John Baptist Gauci), isci1@um.edu.mt (Irene Sciriha)

a *Singular Deck*). This family of graphs has remarkable properties as explored in [2] and [8].

It is the purpose of this work to determine the existence of complete NSSDs. In [8], a characterisation of a graph G which is $\mathbf{G}$ -nutful is given. The inverse $\mathbf{G}^{-1}$ of a $\mathbf{G}$ -nutful graph is associated with a complete NSSD. If $\mathbf{G}$ is the matrix of a complete NSSD graph K_n on n vertices, then the graph of $\mathbf{G}^{-1}$ is said to be $\mathbf{G}^{-1}$ -nutful.

As noted in [2, 4, 7], apart from K_2 , no other graph was known to be $\mathbf{G}$ -nutful. Are there any $\mathbf{G}$ -nutful graphs other than K_2 ? In this paper, we establish that there exist $\mathbf{G}$ -nutful graphs of any order at least six. Some of these $\mathbf{G}$ -nutful graphs turn out to be also complete, in which case both G and the graph of $\mathbf{G}^{-1}$ are nutful.

2 Preliminaries

The *characteristic polynomial* $\phi(\mathbf{M}, \lambda)$ of a square $n \times n$ matrix $\mathbf{M}$, over a field F , is the determinant $\det(\lambda \mathbf{I} - \mathbf{M})$, denoted also by $|\lambda \mathbf{I} - \mathbf{M}|$, which is a polynomial of degree n in λ , where $\mathbf{I}$ denotes the identity matrix. The roots of $\phi(\mathbf{M}, \lambda)$ (in the algebraic closure of F) are the *eigenvalues* of $\mathbf{M}$. For a real and symmetric $n \times n$ matrix $\mathbf{M}$, the n eigenvalues are real, but not necessarily all distinct. The multiplicity of the eigenvalue 0 of $\mathbf{M}$ is called the *nullity* $\eta(\mathbf{M})$ of $\mathbf{M}$. The matrix $\mathbf{M}$ is said to be *singular* if $\eta(\mathbf{M}) > 0$ and is said to be *non-singular* otherwise.

A graph G is a collection of points (called *vertices*) and lines (called *edges*) connecting pairs of vertices. If e is an edge connecting two distinct vertices u and v , then we write $e = \{u, v\}$ and say that e is *incident* to u and to v , or, equivalently, that u and v are *adjacent*. The number of edges incident to a vertex is called the *degree* of the vertex. In a ρ -regular graph, each vertex has degree ρ . For each edge e of G , we associate a non-zero real number w , called the *weight* of e . For a real and symmetric matrix $\mathbf{M}$ whose entries on the leading diagonal are all zero, we can associate a graph $G = \Gamma(\mathbf{M})$ with no loops and with weighted edges, such that $\{i, j\}$ is an edge of G with weight w if and only if the ij^{th} entry of $\mathbf{M}$ is $w \neq 0$. We say that $\mathbf{M}$ is an adjacency matrix representing the graph G , or, equivalently, that G is the graph associated with the adjacency matrix $\mathbf{M}$. The complete graph K_n on n vertices and having $\binom{n}{2}$ weighted edges is represented by the matrix $\mathbf{K}_n$. Although in the study of molecular structures, graphs usually have a maximum vertex degree of at most three, in this study we place no restriction on the maximum vertex degree. This allows us to consider complete graphs on any number of vertices.

In the sequel, we shall also consider vertex-deleted subgraphs of a graph and their associated adjacency matrices. For any vertex x of a graph G , the subgraph $G - x$ obtained by deleting the vertex x and all the edges of G incident to x is associated with the $(n - 1) \times (n - 1)$ matrix $\mathbf{G} - \mathbf{x}$ obtained by deleting the x^{th} row and column of the adjacency matrix $\mathbf{G}$ representing G .

An identity due to Jacobi [6] describes the possible changes in the nullity of an adjacency matrix $\mathbf{G}$ representing the graph G when any two vertices x and y are deleted from G . Denoting the adjugate of $\mathbf{G}$ by $\text{adj } \mathbf{G}$ and the entry in the x, y position of $\text{adj } \mathbf{G}$ by $(\text{adj } \mathbf{G})_{xy}$, we have the following result.

Lemma 2.1. *Let x and y be two distinct vertices of a graph G with associated adjacency matrix $\mathbf{G}$, and let $j = (\text{adj } (\lambda \mathbf{I} - \mathbf{G}))_{xy}$ be the entry in the x, y position of $\text{adj } (\lambda \mathbf{I} - \mathbf{G})$. Then $j^2 = ut - sv$, where s, t, u and v represent the characteristic polynomials of $\mathbf{G}$, $\mathbf{G} - \mathbf{x}$, $\mathbf{G} - \mathbf{y}$ and $\mathbf{G} - \mathbf{x} - \mathbf{y}$, respectively.*

In [2] and [8], the authors introduced a class of graphs termed NSSDs and analysed the properties of these graphs. Formally, NSSDs are defined as follows.

Definition 2.2 ([2]). A graph $G = \Gamma(\mathbf{G})$ is a *NSSD* if $\mathbf{G}$ is a non-singular, real and symmetric matrix with each entry of the diagonal equal to zero, and such that all the $(n - 1) \times (n - 1)$ principal submatrices of $\mathbf{G}$ are singular.

The following is a necessary and sufficient condition for $\Gamma(\mathbf{G})$ to be a NSSD.

Theorem 2.3 ([2]). *The matrices $\mathbf{G}$ and $\mathbf{G}^{-1}$ are real and symmetric with each entry on the respective diagonals equal to zero if and only if $\Gamma(\mathbf{G})$ and $\Gamma(\mathbf{G}^{-1})$ are both NSSDs.*

Therefore, this class of graphs is closed under taking the inverse of the adjacency matrix, hence giving rise to a duality relationship between $\mathbf{G}$ and its inverse $\mathbf{G}^{-1}$, so that either one of the two can assume the principal role. For simplicity, we may refer to $\Gamma(\mathbf{G}^{-1})$ as the inverse of the graph G .

If the only zero entries in a $n \times n$ symmetric matrix are situated on the leading diagonal, then the graph associated with such a matrix is the complete graph K_n . Of special interest to us for this work are complete graphs which also happen to be NSSDs, referred to in the sequel as *complete NSSDs*. In Section 3 we show that the only complete NSSD on at most five vertices is K_2 . In Section 4 we extend the investigation to graphs on at least six vertices, and prove that we can find NSSDs on n vertices for any $n \geq 6$. In particular, the NSSDs we present are complete NSSDs with appropriate edge-weights. In our construction of these complete NSSDs we make use of spectral properties of circulant matrices.

Definition 2.4. A *circulant matrix* is an $n \times n$ matrix $\mathbf{C} = (a_{k,j} : k, j = 1, 2, \dots, n)$ where $a_{k,j} = a_{j-k \pmod n}$, that is

$$\mathbf{C} = \begin{pmatrix} a_0 & a_1 & \cdots & a_{n-2} & a_{n-1} \\ a_{n-1} & a_0 & a_1 & & a_{n-2} \\ \vdots & a_{n-1} & a_0 & \ddots & \vdots \\ a_2 & & \ddots & \ddots & a_1 \\ a_1 & a_2 & \cdots & a_{n-1} & a_0 \end{pmatrix},$$

denoted by $\mathbf{C} = \langle a_0, a_1, \dots, a_{n-1} \rangle$.

3 Complete Graphs On Less Than 6 Vertices

We treat the two cases for $n \leq 4$ and $n = 5$ separately.

3.1 Complete Graphs On Less Than 5 Vertices

For a graph **not** to be a NSSD for any associated adjacency matrix, it suffices to show that it has a non-singular vertex-deleted subgraph. This is shown to be the case for $n = 3$ and 4.

Proposition 3.1. *The complete graph K_n on $n \leq 4$ vertices is a NSSD if and only if $n = 2$.*

Proof. Clearly K_1 is not a NSSD since $\mathbf{K}_1$ is singular. The adjacency matrix representing K_2 is $\begin{pmatrix} 0 & c \\ c & 0 \end{pmatrix}$ for some nonzero real number c . Thus K_2 is a NSSD for any edge weight

c by Definition 2.2. The vertex-deleted subgraphs of K_3 are associated with the non-singular matrices $\mathbf{K}_2$, and hence K_3 cannot be a NSSD for any edge weights. Moreover, the adjacency matrix representing K_3 is $\begin{pmatrix} 0 & a_1 & a_2 \\ a_1 & 0 & a_3 \\ a_2 & a_3 & 0 \end{pmatrix}$ for some nonzero real numbers a_1, a_2, a_3 , whose determinant is $2a_1a_2a_3 \neq 0$. This means that $\mathbf{K}_3$, which represents every vertex-deleted subgraph of K_4 , is always non-singular, implying that K_4 cannot be a NSSD. $\square$

3.2 Complete Graphs On 5 Vertices

Since the weights of K_4 can be assigned such that it can either be singular or non-singular, we cannot utilise the argument used in Proposition 3.1 to prove that K_5 is not a NSSD.

We quote a result from [2] that characterises the nullity of a two-vertex-deleted subgraph of a NSSD depending on whether the two vertices form an edge in its inverse.

Theorem 3.2 ([2]). *If $\Gamma(\mathbf{G})$ is a NSSD on at least three vertices, then $\mathbf{G} - \mathbf{x} - \mathbf{y}$ has nullity zero if $\{x, y\}$ is an edge in $\Gamma(\mathbf{G}^{-1})$, and has nullity two if $\{x, y\}$ is not an edge in $\Gamma(\mathbf{G}^{-1})$.*

With the aim to show that K_5 cannot be a NSSD, we seek properties that a NSSD on five vertices could have.

Lemma 3.3. *If a NSSD on five vertices exists, then it must be the complete graph K_5 for appropriate edge weights, and its inverse must also be K_5 , possibly with different edge weights.*

Proof. Suppose G is a NSSD on five vertices. Then, for any two vertices x and y , $\mathbf{G} - \mathbf{x} - \mathbf{y}$ has nullity zero or two by Theorem 3.2. But $\mathbf{G} - \mathbf{x} - \mathbf{y}$ has only three vertices. There are only four graphs on three vertices, none of which has nullity two, no matter what weights are assigned to their edges. It follows that $\eta(\mathbf{G} - \mathbf{x} - \mathbf{y}) = 0$, and by Theorem 3.2, $\{x, y\}$ is an edge in $\Gamma(\mathbf{G}^{-1})$ for all vertices x and y . Thus, $\Gamma(\mathbf{G}^{-1})$ is K_5 for some particular edge weights. Since $\mathbf{G}^{-1}$ represents a NSSD on five vertices, we repeat the same argument and obtain that its inverse, which is the original matrix $\mathbf{G}$, also represents a complete graph. $\square$

It follows that if a NSSD on five vertices were to exist, it must be complete. Now we make use of Lemma 3.3 to show that K_5 cannot be a NSSD.

Theorem 3.4. *There are no NSSDs on five vertices.*

Proof. By Lemma 3.3, if a NSSD on 5 vertices exists, it must be the graph K_5 with some edge weights, and its inverse (which is also a NSSD on 5 vertices) must also be K_5 , with possibly different edge weights. Let

$$\mathbf{G} = c\mathbf{M} = c \begin{pmatrix} 0 & g_0 & g_1 & g_2 & g_3 \\ g_0 & 0 & g_4 & g_5 & g_6 \\ g_1 & g_4 & 0 & g_7 & g_8 \\ g_2 & g_5 & g_7 & 0 & g_9 \\ g_3 & g_6 & g_8 & g_9 & 0 \end{pmatrix},$$

where $g_i \neq 0$ for all i and $c = \sqrt[5]{|\mathbf{G}|}$, so that $|\mathbf{M}| = 1$. Since $\mathbf{G}$ is a NSSD, so is $\mathbf{M}$. The NSSD

$$\mathbf{G}^{-1} = \frac{1}{c} \mathbf{M}^{-1} = \frac{1}{c} \begin{pmatrix} 0 & h_0 & h_1 & h_2 & h_3 \\ h_0 & 0 & h_4 & h_5 & h_6 \\ h_1 & h_4 & 0 & h_7 & h_8 \\ h_2 & h_5 & h_7 & 0 & h_9 \\ h_3 & h_6 & h_8 & h_9 & 0 \end{pmatrix},$$

where $h_i \neq 0$ for all i . Using Jacobi's Theorem (Lemma 2.1), we have $((\text{adj } \mathbf{M})_{xy})^2 = -|\mathbf{M}||\mathbf{M} - \mathbf{x} - \mathbf{y}| = -|\mathbf{M} - \mathbf{x} - \mathbf{y}|$ for any two vertices x and y of G . But $\mathbf{M} - \mathbf{x} - \mathbf{y} = \mathbf{K}_3$ for all vertices x and y , whose determinant is $2g_i g_j g_k$ for distinct values of $i, j, k \in \{0, \dots, 9\}$. Also $\mathbf{M}_{xy}^{-1} = \frac{(\text{adj } \mathbf{M})_{xy}}{|\mathbf{M}|} = (\text{adj } \mathbf{M})_{xy}$, and thus $h_\ell^2 = -2g_i g_j g_k$ for some i, j, k, ℓ . Specifically:

$$\begin{aligned} -2g_0 g_1 g_4 &= h_9^2 & -2g_0 g_2 g_5 &= h_8^2 & -2g_0 g_3 g_6 &= h_7^2 & -2g_1 g_2 g_7 &= h_6^2 \\ -2g_1 g_3 g_8 &= h_5^2 & -2g_2 g_3 g_9 &= h_4^2 & -2g_4 g_5 g_7 &= h_3^2 & -2g_4 g_6 g_8 &= h_2^2 \\ & & -2g_5 g_6 g_9 &= h_1^2 & -2g_7 g_8 g_9 &= h_0^2 \end{aligned}$$

Considering $\mathbf{M}^{-1}$ as the NSSD, we have:

$$\begin{aligned} -2h_0 h_1 h_4 &= g_9^2 & -2h_0 h_2 h_5 &= g_8^2 & -2h_0 h_3 h_6 &= g_7^2 & -2h_1 h_2 h_7 &= g_6^2 \\ -2h_1 h_3 h_8 &= g_5^2 & -2h_2 h_3 h_9 &= g_4^2 & -2h_4 h_5 h_7 &= g_3^2 & -2h_4 h_6 h_8 &= g_2^2 \\ & & -2h_5 h_6 h_9 &= g_1^2 & -2h_7 h_8 h_9 &= g_0^2 \end{aligned}$$

Squaring the last equation, we have $4h_7^2 h_8^2 h_9^2 = g_0^4$, and multiplying the first three equations together, we obtain $-8g_0^3 g_1 g_2 g_3 g_4 g_5 g_6 = h_7^2 h_8^2 h_9^2$. Thus $-32g_1 g_2 g_3 g_4 g_5 g_6 = g_0$. Since $-2g_7 g_8 g_9 = h_0^2$, we can write

$$16g_0 g_1 g_2 g_3 g_4 g_5 g_6 g_7 g_8 g_9 = g_0^2 h_0^2$$

This process can be repeated starting from another group of equations to obtain $16 \prod_{j=0}^9 g_j = g_i^2 h_i^2$ for all $i \in \{0, 1, \dots, 9\}$. Thus $g_i^2 h_i^2 = g_j^2 h_j^2 = 16 \prod_{k=0}^9 g_k$ for all $i, j \in \{0, 1, \dots, 9\}$.

Since $\mathbf{M}\mathbf{M}^{-1} = \mathbf{I}$, we have, in particular, that $g_0 h_0 + g_1 h_1 + g_2 h_2 + g_3 h_3 = 1$. But $g_i h_i = \pm g_j h_j$ for all i, j , and hence $g_0 h_0 (\pm 1 \pm 1 \pm 1 \pm 1) = 1$. Thus $g_0 h_0 = \pm \frac{1}{4}$ or $g_0 h_0 = \pm \frac{1}{2}$. Since $g_i^2 h_i^2 = g_j^2 h_j^2$ for all i, j then either $g_i h_i = \pm \frac{1}{4}$ for all i or $g_i h_i = \pm \frac{1}{2}$ for all i . Additionally, in order for $g_0 h_0 + g_1 h_1 + g_2 h_2 + g_3 h_3$ to be equal to 1, none of the $g_i h_i$'s can be equal to $-\frac{1}{4}$, and thus either all $g_i h_i$ are equal to $\frac{1}{4}$ or all $g_i h_i$ are equal to $\pm \frac{1}{2}$.

Suppose $g_i h_i = \pm \frac{1}{2}$ for all i . In the five equations

$$\begin{aligned} g_0 h_0 + g_1 h_1 + g_2 h_2 + g_3 h_3 &= 1 \\ g_0 h_0 + g_4 h_4 + g_5 h_5 + g_6 h_6 &= 1 \\ g_1 h_1 + g_4 h_4 + g_7 h_7 + g_8 h_8 &= 1 \\ g_2 h_2 + g_5 h_5 + g_7 h_7 + g_9 h_9 &= 1 \\ g_3 h_3 + g_6 h_6 + g_8 h_8 + g_9 h_9 &= 1, \end{aligned}$$

three of the four terms $g_i h_i$ in each equation must have value $\frac{1}{2}$ and the remaining one must be equal to $-\frac{1}{2}$. Combinatorially, this is impossible, for suppose we choose, without loss of generality, $g_0 h_0 = -\frac{1}{2}$; then $g_1 h_1 = g_2 h_2 = g_3 h_3 = g_4 h_4 = g_5 h_5 = g_6 h_6 = \frac{1}{2}$.

From the last three equations, we obtain that $g_7h_7 + g_8h_8 = 0$, $g_7h_7 + g_9h_9 = 0$ and $g_8h_8 + g_9h_9 = 0$, implying that $g_7h_7 = g_8h_8 = g_9h_9 = 0$, a contradiction.

The only remaining possibility is that $g_ih_i = \frac{1}{4}$ for all i . Consider $\mathbf{M}\mathbf{M}^{-1} = \mathbf{M}^{-1}\mathbf{M} = \mathbf{I}$. In particular, by multiplying the first row of $\mathbf{M}$ by the second column of $\mathbf{M}^{-1}$, and by multiplying the first row of $\mathbf{M}^{-1}$ by the second column of $\mathbf{M}$, we obtain

$$\begin{aligned} g_1h_4 + g_2h_5 + g_3h_6 &= 0 \\ h_1g_4 + h_2g_5 + h_3g_6 &= 0 \end{aligned}$$

Since $g_ih_i = \frac{1}{4}$, we can write

$$\begin{aligned} \frac{1}{4} \left(\frac{g_1}{g_4} + \frac{g_2}{g_5} + \frac{g_3}{g_6} \right) &= 0 \\ \frac{1}{4} \left(\frac{g_4}{g_1} + \frac{g_5}{g_2} + \frac{g_6}{g_3} \right) &= 0 \end{aligned}$$

Let $p = \frac{g_1}{g_4}$, $q = \frac{g_2}{g_5}$ and $r = \frac{g_3}{g_6}$ so that

$$p + q + r = 0 \quad (3.1)$$

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 0 \quad (3.2)$$

From (3.2) we obtain $pq + pr + qr = 0$. Substituting (3.1) in this equation, we get $(-q - r)q + (-q - r)(r) + qr = 0$, or $q^2 + qr + r^2 = 0$. Fixing r to be any real number, this is a quadratic equation in q with a negative discriminant $-3r^2$, and hence q must be a complex number, which is a contradiction.

Thus $\Gamma(\mathbf{M})$ is not a NSSD, and hence neither is $\Gamma(\mathbf{G})$. $\square$

From Proposition 3.1 and Theorem 3.4, the following result follows immediately.

Corollary 3.5. *The complete graph K_n on $n \leq 5$ vertices is a NSSD if and only if $n = 2$.*

4 Complete Graphs on at Least Six Vertices

In this section, we prove that there indeed exist complete NSSDs on at least six vertices by a direct construction.

Any complete graph on $n \geq 6$ vertices has at least one (Hamiltonian) cycle of length n on all the vertices. If we assign an arbitrary weight $a \neq 0$ to this cycle and a weight of 1 to every other edge of the graph, the matrix obtained is $\mathbf{K}_n^{(a)}$ of the form shown in (4.1):

$$\mathbf{K}_n^{(a)} = \begin{pmatrix} 0 & a & 1 & \cdots & 1 & a \\ a & 0 & a & 1 & \cdots & 1 \\ 1 & a & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & a & 1 \\ 1 & \cdots & 1 & a & 0 & a \\ a & 1 & \cdots & 1 & a & 0 \end{pmatrix} \quad (4.1)$$

4.1 The Characteristic Polynomial of Circulant Matrices

We remark that the real and symmetric matrix shown in (4.1) is the circulant matrix $\langle 0, a, 1, \dots, 1, a \rangle$. The eigenvalues of such a matrix can thus be obtained by using well-known results on circulant matrices.

Lemma 4.1. *The eigenvalues of the circulant matrix $\langle 0, a, 1, \dots, 1, a \rangle$ are $\lambda_0 = 2(a - 1) - 1 + n$, and $\lambda_j = 2(a - 1) \cos\left(\frac{2\pi j}{n}\right) - 1$ for $j \in \{1, \dots, n-1\}$, $a \neq 0$.*

Proof. It is well-known (see, for example, [5]) that the eigenvalues of the general circulant matrix $\langle a_0, a_1, \dots, a_{n-1} \rangle$ are $\lambda_j = \sum_{k=0}^{n-1} a_k \omega_j^k$, for $j \in \{0, \dots, n-1\}$ and $\omega_0, \dots, \omega_{n-1}$ are the n^{th} roots of unity given by $\omega_j = e^{2\pi i j/n}$. If we substitute $a_0 = 0$, $a_2 = a_3 = \dots = a_{n-2} = 1$ and $a_{n-1} = a_1 = a$, then the eigenvalues of $\langle 0, a, 1, \dots, 1, a \rangle$ are given by $\lambda_j = a(\omega_j + \omega_j^{n-1}) + \sum_{k=2}^{n-2} \omega_j^k$. But

$$\sum_{k=0}^{n-1} \omega_j^k = \begin{cases} \sum_{k=0}^{n-1} 1^k = n & \text{if } j = 0 \\ \sum_{k=0}^{n-1} (-1)^k = 0 & \text{if } n \text{ is even and } j = \frac{n}{2} \\ \frac{(\omega_j^n)^n - 1}{\omega_j^n - 1} = \frac{(\omega_j^n)^k - 1}{\omega_j^k - 1} = 0 & \text{otherwise} \end{cases}$$

Hence, if $j = 0$, then $\sum_{k=2}^{n-2} \omega_0^k = n - 3$, and thus

$$\lambda_0 = a(1 + 1) + (n - 3) = 2(a - 1) - 1 + n.$$

If n is even and $j = \frac{n}{2}$, then $\sum_{k=2}^{n-2} \omega_{n/2}^k = 0 - 1 + 1 + 1 = 1$. Thus

$$\begin{aligned} \lambda_{n/2} &= a((-1) + (-1)^{n-1}) + 1 = a((-1)(1 + (-1)^{n-2}) + 1) = 2a(\cos \pi) + 1 \\ &= 2(a - 1)(\cos \pi) - 1 = 2(a - 1) \cos\left(\frac{2\pi j}{n}\right) - 1. \end{aligned}$$

For any other case, $\sum_{k=2}^{n-2} \omega_j^k = -1 - \omega_j - \omega_j^{n-1}$, which means that

$$\lambda_j = (a - 1)(\omega_j + \omega_j^{n-1}) - 1 = (a - 1)(\omega_j + \omega_j^{-1}) - 1 = 2(a - 1) \cos\left(\frac{2\pi j}{n}\right) - 1,$$

completing the proof. $\square$

The eigenvalues of $\langle 0, a, 1, \dots, 1, a \rangle$ satisfy several interesting properties. To order the eigenvalues, we use the notation of Lemma 4.1.

Proposition 4.2. Let $n \geq 6$ and let the complete graph K_n have circulant matrix $\mathbf{K}_n^{(a)}$ given by $\mathbf{K}_n^{(a)} = \langle 0, a, 1, \dots, 1, a \rangle$, where the parameter a is a non-zero real number. Then the eigenvalues $\lambda_0, \dots, \lambda_{n-1}$ of $\mathbf{K}_n^{(a)}$ are such that

(i) $\lambda_0 > \lambda_j$ for all $j = 1, \dots, n-1$;

(ii) $\lambda_j = \lambda_{n-j}$ for $j = 1, \dots, \lfloor \frac{n-1}{2} \rfloor$;

(iii) $\lambda_j > \lambda_{j+1}$ ($j = 1, \dots, \lfloor \frac{n-1}{2} \rfloor$) when $a > 1$.

For $\mathbf{K}_8^{(2)}$, say, $\lambda_0 = 15$, $\lambda_1 = \lambda_7 = -1 + 4\sqrt{2}$, $\lambda_2 = \lambda_6 = -1$, $\lambda_3 = \lambda_5 = -1 - 4\sqrt{2}$ and $\lambda_4 = -9$.

The characteristic polynomial of $\mathbf{K}_n^{(a)} = \langle 0, a, 1, \dots, 1, a \rangle$ can be obtained from Lemma 4.1. However, we can actually also determine the characteristic polynomials of all the vertex-deleted subgraphs of $\mathbf{K}_n^{(a)}$. In the sequel, the characteristic polynomial $\phi(\mathbf{K}_n^{(a)}, \lambda)$ will be denoted by $\psi(\mathbf{K}_n^{(a)}, x, c)$ where $x = \lambda$ and $c = 2(a-1)$, $c \neq -2$.

Proposition 4.3. Let n be any integer greater than 1 and let $\mathbf{K}_n^{(a)}$ be a complete circulant symmetric matrix $\langle 0, a, 1, \dots, 1, a \rangle$ on n vertices, $a \neq 0$. Then its characteristic polynomial is

$$\psi(\mathbf{K}_n^{(a)}, x, c) = \begin{cases} (x+1-c-n)(p(x, c))^2 & \text{if } n \text{ is odd} \\ (x+1-c-n)(x+1+c)(p(x, c))^2 & \text{if } n \text{ is even} \end{cases}$$

and $\psi(\mathbf{K}_n^{(a)} - \mathbf{v}, x, c)$, the characteristic polynomial of each of its vertex-deleted subgraphs, is

$$\begin{cases} \frac{p(x, c)}{n} (p(x, c) + 2(x+1-c-n)q(x, c)) & \text{if } n \text{ is odd} \\ \frac{p(x, c)}{n} ((2x+2-n)p(x, c) + 2(x+1-c-n)(x+1+c)q(x, c)) & \text{if } n \text{ is even} \end{cases}$$

where $k = \lfloor \frac{n-1}{2} \rfloor$, $c = 2(a-1)$, $p(x, c) = \prod_{j=1}^k (x+1-c \cos(\frac{2\pi j}{n}))$ and

$$q(x, c) = \frac{\partial p}{\partial x} = \sum_{j=1}^k \prod_{q=1, q \neq j}^k (x+1-c \cos(\frac{2\pi q}{n})).$$

Proof. By Lemma 4.1, the eigenvalues of $\mathbf{K}_n^{(a)}$ are $\lambda_0 = 2(a-1) - 1 + n$ and $\lambda_j = 2(a-1) \cos(\frac{2\pi j}{n}) - 1$ for $j \in \{1, \dots, n-1\}$. Furthermore, by Proposition 4.2 (ii), $\lambda_{n-j} = \lambda_j$ for all $j \in \{1, \dots, k\}$. Also, if n is even, then $\lambda_{n/2} = -2(a-1) - 1$. Hence, for $c = 2(a-1)$, the characteristic polynomial of $\mathbf{K}_n^{(a)}$ is

$$\prod_{j=0}^{n-1} (x - \lambda_j) = (x+1-c-n) \prod_{j=1}^k (x - \lambda_j)^2 = (x+1-c-n) \left(\prod_{j=1}^k (x+1-c \cos(\frac{2\pi j}{n})) \right)^2$$

when n is odd, and

$$\begin{aligned} \prod_{j=0}^{n-1} (x - \lambda_j) &= (x + 1 - c - n)(x + 1 + c) \prod_{j=1}^k (x - \lambda_j)^2 \\ &= (x + 1 - c - n)(x + 1 + c) \left(\prod_{j=1}^k \left(x + 1 - c \cos \left(\frac{2\pi j}{n} \right) \right) \right)^2 \end{aligned}$$

when n is even.

Since every circulant matrix associated with a graph G renders G vertex-transitive, all the vertex-deleted subgraphs of G are isomorphic and hence have the same characteristic polynomial. But the sum of the characteristic polynomials of all the vertex-deleted subgraphs of G is equal to the derivative of the characteristic polynomial of G [9]¹. Thus $\psi(\mathbf{K}_n^{(a)} - \mathbf{v}, x, c) = \frac{1}{n} \frac{\partial}{\partial x} \psi(\mathbf{K}_n^{(a)}, x, c)$, so we now proceed to evaluate this partial derivative.

When n is odd, we have

$$\begin{aligned} \psi(\mathbf{K}_n^{(a)} - \mathbf{v}, x, c) &= \frac{1}{n} \frac{\partial}{\partial x} (x + 1 - c - n)(p(x, c))^2 \\ &= \frac{1}{n} \left((p(x, c))^2 + 2(x + 1 - c - n)p(x, c) \frac{\partial p}{\partial x} \right) \\ &= \frac{p(x, c)}{n} \left(p(x, c) + 2(x + 1 - c - n) \left(\frac{\partial}{\partial x} \prod_{j=1}^k (x + 1 - c \cos \left(\frac{2\pi j}{n} \right)) \right) \right) \\ \psi(\mathbf{K}_n^{(a)} - \mathbf{v}, x, c) &= \frac{p(x, c)}{n} \left(p(x, c) + 2(x + 1 - c - n) \sum_{j=1}^k \prod_{q=1, q \neq j}^k (x + 1 - c \cos \left(\frac{2\pi q}{n} \right)) \right) \end{aligned}$$

When n is even, the result stated in the proposition statement is obtained in a similar manner. □

4.2 The Construction of Complete NSSDs

We have noted in the proof of Proposition 4.3 that the characteristic polynomials of the adjacency matrices of all the vertex-deleted subgraphs of the graph associated with $\mathbf{K}_n^{(a)} = \langle 0, a, 1, \dots, 1, a \rangle$ are equal. This means that the determinant of the matrix of each of these vertex-deleted subgraphs is the same. Thus, if the determinant of $\mathbf{K}_n^{(a)}$ is non-zero and the determinant of any one of its $(n-1) \times (n-1)$ principal submatrices is zero, then the matrix would represent a NSSD. This is the basis of our construction.

To illustrate the above, let us take the complete graph on seven vertices as an example. By Corollary 3.5 this is the smallest odd number of vertices we can take to possibly find a

¹This is a well-known result for graphs with 0–1 adjacency matrices. However, this result has been known to hold for any graph with a real and symmetric adjacency matrix from as early as 1968.

NSSD. The matrix $\mathbf{K}_7^{(a)}$ would be of the form

$$\mathbf{K}_7^{(a)} = \begin{pmatrix} 0 & a & 1 & 1 & 1 & 1 & a \\ a & 0 & a & 1 & 1 & 1 & 1 \\ 1 & a & 0 & a & 1 & 1 & 1 \\ 1 & 1 & a & 0 & a & 1 & 1 \\ 1 & 1 & 1 & a & 0 & a & 1 \\ 1 & 1 & 1 & 1 & a & 0 & a \\ a & 1 & 1 & 1 & 1 & a & 0 \end{pmatrix}$$

which has determinant

$$\det(\mathbf{K}_7^{(a)}) = 2(2+a)(f(a))^2$$

where

$$f(a) = -1 + 2a + a^2 - a^3$$

The determinant of each $(n-1) \times (n-1)$ principal submatrix of $\mathbf{K}_7^{(a)}$ is

$$\det(\mathbf{K}_7^{(a)} - \mathbf{v}) = f(a)(1 - 6a - a^2 + a^3)$$

We require a value of a such that $\det(\mathbf{K}_7^{(a)}) \neq 0$ and $\det(\mathbf{K}_7^{(a)} - \mathbf{v}) = 0$. We have the conditions:

$$a \neq 0 \quad (4.2)$$

$$2 + a \neq 0 \quad (4.3)$$

$$f(a) \neq 0 \quad (4.4)$$

$$1 - 6a - a^2 + a^3 = 0 \quad (4.5)$$

Thus if we find a value for a that satisfies (4.2), (4.3), (4.4) and (4.5), then a complete NSSD is obtained. We note that the cubic equation (4.5) has the real solutions $a = -2.0938, 0.16296, 2.9308$, which also satisfy (4.2), (4.3) and (4.4). Indeed, if we take $a = 2.9308$, then the inverse of $\mathbf{K}_7^{(a)}$ is

$$\left(\mathbf{K}_7^{(a)}\right)^{-1} = \begin{pmatrix} 0 & 0.2327 & 0.068 & -0.25 & -0.25 & 0.068 & 0.2327 \\ 0.2327 & 0 & 0.2327 & 0.068 & -0.25 & -0.25 & 0.068 \\ 0.068 & 0.2327 & 0 & 0.2327 & 0.068 & -0.25 & -0.25 \\ -0.25 & 0.068 & 0.2327 & 0 & 0.2327 & 0.068 & -0.25 \\ -0.25 & -0.25 & 0.068 & 0.2327 & 0 & 0.2327 & 0.068 \\ 0.068 & -0.25 & -0.25 & 0.068 & 0.2327 & 0 & 0.2327 \\ 0.2327 & 0.068 & -0.25 & -0.25 & 0.068 & 0.2327 & 0 \end{pmatrix}$$

which is the circulant symmetric matrix $\langle 0, 0.2327, 0.068, -0.25, -0.25, 0.068, 0.2327 \rangle$. For $a = -2.0938$ or 0.16296 , a similar result is obtained.

We remark that the inverse of a non-singular circulant matrix is another circulant matrix; this is evident in $\left(\mathbf{K}_7^{(a)}\right)^{-1}$ above. Furthermore, $\left(\mathbf{K}_7^{(a)}\right)^{-1}$ represents a complete graph on seven vertices such that each one of its three edge-disjoint cycles has edges of weight $0.2327, 0.068$ and -0.25 , respectively.

Using a similar technique, we can show that if $\mathbf{K}_6^{(a)} = \langle 0, a, 1, 1, 1, a \rangle$, then $a = -1 \pm \sqrt{2}$ for $\mathbf{K}_6^{(a)}$ to be a NSSD. Indeed, if we choose $a = -1 - \sqrt{2} \approx -2.4142$, then

$(\mathbf{K}_6^{(a)})^{-1} = \langle 0, -0.6306, 1.5224, -1.5224, 1.5224, -0.6306 \rangle$. Furthermore, if $\mathbf{K}_8^{(a)} = \langle 0, a, 1, 1, 1, 1, 1, a \rangle$, then, using the above method, we can show that when $a = 2.1889$, $(\mathbf{K}_8^{(a)})^{-1} = \langle 0, -0.2592, 0.3459, 0.1898, -0.3082, 0.1898, 0.3459, -0.2592 \rangle$. Hence, we have:

Theorem 4.4. *There exist complete NSSDs on 6, 7 and 8 vertices.*

4.3 Existence of Complete NSSDs on at least Nine Vertices

The method described in the previous subsection may go wrong in the case that a polynomial equation $\xi(a) = 0$ akin to (4.5) having only complex solutions is obtained. Indeed, this scenario happens if we apply our construction to $\mathbf{K}_5^{(a)}$; in this case the quadratic polynomial $\xi(a)$ is $a^2 + a + 1$, which has complex roots. Thus, there is no guarantee that the above construction will always yield a complete NSSD unless we prove that $\xi(a)$ has always at least one real root. This is what we do in the next theorem.

Theorem 4.5. *Let n be any integer greater than 8 and let $\mathbf{K}_n^{(a)}$ be a complete circulant symmetric matrix $\langle 0, a, 1, \dots, 1, a \rangle$ on n vertices, $a \neq 0$. Then there exists a value of the parameter a strictly between 1 and $1 + \frac{1}{2} \sec\left(\frac{4\pi}{n}\right)$ such that K_n is a NSSD.*

Proof. We can determine the characteristic polynomials of $\mathbf{K}_n$ and $\mathbf{K}_n - \mathbf{v}$ from Proposition 4.3. For K_n to be a NSSD, we need to find a value of c such that $\psi(\mathbf{K}_n^{(a)}, 0, c) \neq 0$ and $\psi(\mathbf{K}_n^{(a)} - \mathbf{v}, 0, c) = 0$, recalling that $c = 2(a - 1)$. Considering first the case when n is an odd number greater than 8, we obtain:

$$\begin{aligned} (1 - c - n)(p(0, c))^2 &\neq 0 \\ \frac{p(0, c)}{n} \left(p(0, c) + 2(1 - c - n) \sum_{j=1}^k \prod_{q=1, q \neq j}^k (1 - c \cos\left(\frac{2\pi q}{n}\right)) \right) &= 0 \end{aligned}$$

where $k = \frac{n-1}{2}$ and $p(0, c)$ is as defined in Proposition 4.3. Hence $c \neq 1 - n$, $p(0, c) \neq 0$ and $h(c) = 0$, where

$$h(c) = \prod_{j=1}^k (1 - c \cos\left(\frac{2\pi j}{n}\right)) + 2(1 - c - n) \sum_{j=1}^k \prod_{q=1, q \neq j}^k (1 - c \cos\left(\frac{2\pi q}{n}\right)) \quad (4.6)$$

We note that $h(c)$ in (4.6) is a polynomial in c with real coefficients and hence it is continuous. Thus, if we show that there exist two distinct values of c for which $h(c)$ has opposite signs, then (4.6) has a real root by the intermediate value theorem.

$$\begin{aligned} \text{Substituting } c = 0 \text{ in (4.6), we obtain } h(0) &= \left(\prod_{j=1}^k 1 \right) + 2(1 - n) \sum_{j=1}^k \left(\prod_{q=1, q \neq j}^k 1 \right) \\ &= 1 + 2(1 - n) \left(\frac{n-1}{2} \right) = 1 - (n-1)^2 < 0 \text{ since } n \geq 9. \end{aligned}$$

We now substitute an appropriate value of c such that $h(c) > 0$. Note, first of all, that

$\sec\left(\frac{4\pi}{n}\right) > 0$ since $n \geq 9$. Substituting this value for c in (4.6), we obtain

$$h\left(\sec\left(\frac{4\pi}{n}\right)\right) = \prod_{j=1}^k \left(1 - \sec\left(\frac{4\pi}{n}\right) \cos\left(\frac{2\pi j}{n}\right)\right) + 2\left(1 - \sec\left(\frac{4\pi}{n}\right) - n\right) \sum_{j=1}^k \prod_{q=1, q \neq j}^k \left(1 - \sec\left(\frac{4\pi}{n}\right) \cos\left(\frac{2\pi q}{n}\right)\right).$$

For $j = 2$, $\left(1 - \sec\left(\frac{4\pi}{n}\right) \cos\left(\frac{2\pi j}{n}\right)\right) = 0$. Thus,

$$h\left(\sec\left(\frac{4\pi}{n}\right)\right) = 2\left(1 - \sec\left(\frac{4\pi}{n}\right) - n\right) \prod_{q=1, q \neq 2}^k \left(1 - \sec\left(\frac{4\pi}{n}\right) \cos\left(\frac{2\pi q}{n}\right)\right)$$

implying that

$$h\left(\sec\left(\frac{4\pi}{n}\right)\right) = 2\left(1 - \sec\left(\frac{4\pi}{n}\right) - n\right) \left(1 - \frac{\cos\left(\frac{2\pi}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}\right) \left(\prod_{q=3}^k \left(1 - \frac{\cos\left(\frac{2\pi q}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}\right)\right). \quad (4.7)$$

We now check the signs of each bracket in (4.7). Since $1 \leq \sec\left(\frac{4\pi}{n}\right) \leq \sec\left(\frac{4\pi}{9}\right) < 6$, we have that $1 - \sec\left(\frac{4\pi}{n}\right) - n < 0$. Thus the first bracket of (4.7) is negative.

The second bracket of (4.7) is

$$1 - \frac{\cos\left(\frac{2\pi}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)} = \frac{\cos\left(\frac{4\pi}{n}\right) - \cos\left(\frac{2\pi}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)} = \frac{-2 \sin\left(\frac{3\pi}{n}\right) \sin\left(\frac{\pi}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}$$

which is negative since $n \geq 9$ and $\sin \theta > 0$ when $0 < \theta < \pi$.

The third bracket of (4.7) is

$$\begin{aligned} \prod_{q=3}^k \left(1 - \frac{\cos\left(\frac{2\pi q}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}\right) &= \prod_{q=3}^k \left(\frac{\cos\left(\frac{4\pi}{n}\right) - \cos\left(\frac{2\pi q}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}\right) \\ &= \prod_{q=3}^k \left(\frac{2 \sin\left(\frac{\pi(q+2)}{n}\right) \sin\left(\frac{\pi(q-2)}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}\right) \end{aligned} \quad (4.8)$$

Since the highest value of q is $k = \frac{n-1}{2}$, then the largest angle in (4.8) is $\frac{\pi\left(\frac{n-1}{2} + 2\right)}{n} = \frac{\pi(n+3)}{2n} < \pi$. Because of this, all the products in (4.8) are positive, and this means that the third bracket of (4.7) is positive. Thus, $h\left(\sec\left(\frac{4\pi}{n}\right)\right)$ in (4.7) is positive.

We now consider the case when n is even. By Proposition 4.3, we obtain

$$\begin{aligned} (1 - c - n)(1 + c)(p(0, c))^2 &\neq 0 \\ \frac{p(0, c)}{n} \left((2 - n)p(0, c) + 2(1 - c - n)(1 + c) \sum_{j=1}^k \prod_{q=1, q \neq j}^k (1 - c \cos\left(\frac{2\pi q}{n}\right)) \right) &= 0 \end{aligned}$$

where $k = \frac{n-2}{2}$. Hence $c \neq 1 - n$, $c \neq -1$, $p(0, c) \neq 0$ and $g(c) = 0$, where

$$g(c) = (2 - n) \prod_{j=1}^k \left(1 - c \cos\left(\frac{2\pi j}{n}\right)\right) + 2(1 - c - n)(1 + c) \sum_{j=1}^k \prod_{q=1, q \neq j}^k \left(1 - c \cos\left(\frac{2\pi q}{n}\right)\right) \quad (4.9)$$

As for the case when n was odd, since $g(c)$ in (4.9) is a polynomial in c , we search for two values of c such that $g(c)$ is positive for one value of c and negative for the other. We show that $c = 0$ and $c = \sec\left(\frac{4\pi}{n}\right)$ are again two such values of c .

Substituting $c = 0$ in (4.9), we obtain

$$g(0) = (2 - n) \left(\prod_{j=1}^k 1 \right) + 2(1 - n)(1) \sum_{j=1}^k \left(\prod_{q=1, q \neq j}^k 1 \right) = (2 - n) + 2(1 - n) \left(\frac{n - 2}{2} \right) = -n(n - 2) < 0 \text{ since } n \geq 9.$$

Substituting $c = \sec\left(\frac{4\pi}{n}\right)$ in (4.9), we obtain

$$\begin{aligned} g\left(\sec\left(\frac{4\pi}{n}\right)\right) &= (2 - n) \prod_{j=1}^k \left(1 - \sec\left(\frac{4\pi}{n}\right) \cos\left(\frac{2\pi j}{n}\right)\right) \\ &\quad + 2\left(1 - \sec\left(\frac{4\pi}{n}\right) - n\right) \left(1 + \sec\left(\frac{4\pi}{n}\right)\right) \sum_{j=1}^k \prod_{q=1, q \neq j}^k \left(1 - \sec\left(\frac{4\pi}{n}\right) \cos\left(\frac{2\pi q}{n}\right)\right) \\ g\left(\sec\left(\frac{4\pi}{n}\right)\right) &= 2\left(1 - \sec\left(\frac{4\pi}{n}\right) - n\right) \left(1 + \sec\left(\frac{4\pi}{n}\right)\right) \prod_{q=1, q \neq 2}^k \left(1 - \sec\left(\frac{4\pi}{n}\right) \cos\left(\frac{2\pi q}{n}\right)\right) \\ g\left(\sec\left(\frac{4\pi}{n}\right)\right) &= 2\left(1 - \sec\left(\frac{4\pi}{n}\right) - n\right) \left(1 + \sec\left(\frac{4\pi}{n}\right)\right) \left(1 - \frac{\cos\left(\frac{2\pi}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}\right) \\ &\quad \left(\prod_{q=3}^k \left(1 - \frac{\cos\left(\frac{2\pi q}{n}\right)}{\cos\left(\frac{4\pi}{n}\right)}\right)\right) \quad (4.10) \end{aligned}$$

Since $1 + \sec\left(\frac{4\pi}{n}\right) > 0$ and the product of all the other brackets in (4.10) was shown to be positive in the proof for the case when n is odd, $n \geq 9$, then $g\left(\sec\left(\frac{4\pi}{n}\right)\right)$ in (4.10) is positive.

Consequently, the polynomials (4.6) and (4.9) both have a root between $c = 0$ and $c = \sec\left(\frac{4\pi}{n}\right)$. Since $c = 2(a - 1)$, the graph associated with the matrix $\mathbf{K}_n^{(a)}$ represents a NSSD for a value of a between 1 and $1 + \frac{1}{2} \sec\left(\frac{4\pi}{n}\right)$, as required. $\square$

Remark 4.6. Observe that the lower bound for the parameter a in Theorem 4.5 can be improved to $\frac{3}{2}$. Since the upper bound tends to $\frac{3}{2}$ as n increases, we know that the adjacency matrix corresponding to a complete NSSD on a large number of vertices is given by $\langle 0, a, 1, \dots, 1, a \rangle$ where $a = 1.5 + \varepsilon$ for a sufficiently small positive real value of ε .

We thus have the following existence result from Theorem 4.4 and Theorem 4.5.

Corollary 4.7. *For each $n \geq 6$, there always exists a complete NSSD on n vertices.*

In [8], a subclass of NSSDs referred to as **G**–nutful graphs was introduced and defined as follows.

Definition 4.8. A **G**–nutful graph G is either K_2 or a NSSD on at least three vertices having all two vertex–deleted subgraphs of the same nullity.

Recall that a necessary and sufficient condition for G to be a **G**–nutful graph is for its inverse to be a complete NSSD. We have established that besides K_2 , there exist other **G**–nutful graphs.

Theorem 4.9. *There exist **G**–nutful graphs on any number of vertices greater than five.*

5 Conclusion

The complete graph K_2 has been well–known to be a complete NSSD. In this paper, we have shown that complete NSSDs exist for all positive integers at least six. This disproves a conjecture that appeared in [8] that NSSDs must have an even number of vertices. Indeed, NSSDs on an odd number vertices other than those that are complete graphs have also been found. An example is given by the graph with circulant adjacency matrix $\langle 0, -0.7549, 1, 0, 0, 1, -0.7549 \rangle$, whose inverse is given by the circulant matrix $\langle 0, 0.1075, 0.5812, 0.3312, 0.3312, 0.5812, 0.1075 \rangle$. However, we have also shown that no complete NSSDs exist on n vertices for $3 \leq n \leq 5$.

In [4, 7] it was conjectured that K_2 is the only *nuciferous graph*, that is a NSSD with 0–1 entries in its adjacency matrix whose inverse is complete. It is still an open problem whether a construction of complete NSSDs similar to that given in Section 4 may lead to an inverse with only 0–1 entries, which would then be nuciferous.

References

- [1] F. Cataldo, A. Graovac and O. Ori (eds.), *The Mathematics and Topology of Fullerenes*, Springer, New York, 2011.
- [2] A. Farrugia, J. B. Gauci and I. Sciriha, On the inverse of the adjacency matrix of a graph, *Special Matrices* **1** (2013), 28–41.
- [3] P. W. Fowler and D. E. Manolopoulos (eds.), *An Atlas of Fullerenes*, Clarendon Press, 1995.
- [4] P. W. Fowler, B. T. Pickup, T. Z. Todorova, R. de los Reyes and I. Sciriha, Omni–conducting fullerenes, *Chemical Physics Letters* (2013), 33–35.
- [5] R. M. Gray, Toeplitz and circulant matrices: a review, *Foundations and Trends in Communications and Information Theory* **2** (2006), 155–239.
- [6] I. Gutman and O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, Berlin, 1986.
- [7] I. Sciriha, M. Debono, M. Borg, P. Fowler and B. T. Pickup, Interlacing–extremal graphs, *Ars Mathematica Contemporanea* (2013), 261–278.
- [8] I. Sciriha, A. Farrugia and J. B. Gauci, The adjacency matrices of complete and nutful graphs, *Match Commun. Math. Comput. Chem* **72** (2014), 165–178.

- [9] R. C. Thompson, Principal submatrices v: some results concerning principal submatrices of arbitrary matrices, *Journal of Research of the National Bureau of Standards—B. Mathematical Sciences* **72B** (1968), 115–125.